

General Compact Quantum Groups, a Tutorial

TOM H. KOORNWINDER

Abstract A tutorial introduction is given to general Hopf algebras and to general compact quantum groups. In the definition and further treatment of compact quantum groups C^* -algebras are avoided. Contact with Woronowicz's compact matrix quantum groups is made at a later stage.

0. Introduction

It is the purpose of this paper to give a tutorial introduction to Hopf algebras and general compact quantum groups. An earlier version of the paper was part of the lecture notes of my course on “Quantum groups and q -special functions” at the European School of Group Theory 1993, Trento, Italy. Eventually, this paper may appear as part of a book containing the notes of the main courses at this School.

For the reader who is new to this subject, it is crucial that he complements reading of the present paper with the study of papers dealing with $SU_q(2)$ and other special quantum groups.

I now describe the various sections in some more detail. Section 1 presents the basic theory of Hopf algebras and of corepresentations. Here (and in §2) proofs are either given in full detail or they are sketched such that the reader can easily fill in the gaps. No attempt has been made to high mathematical sophistication or big generality.

Section 2 deals with compact quantum groups. The original plan was to give here an account of Woronowicz's celebrated theory [9] of compact matrix quantum groups, but this section grew out into an alternative approach to compact quantum groups, avoiding C^* -algebras in the definition and in the proofs, but formulating everything on the Hopf $*$ -algebra level. The C^* -algebra completion now appears as a final observation instead of an essential part of the definition. Both in [9] and in §2 the Haar functional plays a crucial role. The approach of §2, developed joint with M. S. Dijkhuizen [3], may be somewhat shorter and easier to grasp than the C^* -algebra approach. We believe that this approach is very well suited for application to the special compact matrix quantum groups most commonly studied nowadays. The section concludes with a comparison of various approaches to compact quantum groups which have appeared in the literature. Particular mention here deserves the paper [4] by Effros & Ruan, who earlier introduced the same algebras as our CQG

University of Amsterdam, Faculty of Mathematics and Computer Science, Plantage Muidergracht 24, 1018 TV Amsterdam. The Netherlands, email: thk@fwi.uva.nl

algebras, but called them differently and also developed the theory in a different direction.

I thank Mathijs Dijkhuizen for commenting on a preliminary version of this paper.

1. Generalities about Hopf algebras

Standard references about Hopf algebras are the books by Abe [1] and Sweedler [6], see also Hazewinkel [5, §37.1]. Below we will assume ground field $\mathbb{C}$. By the *tensor product* $V \otimes W$ of two linear spaces V and W we will always mean the *algebraic* tensor product. Thus the elements of $V \otimes W$ are finite sums of elements $v_i \otimes w_i$ ($v_i \in V$, $w_i \in W$).

1.1. Hopf algebras. The reader will be familiar with the concept of an *associative algebra with unit* (or shortly an *algebra*), i.e., a linear space $\mathcal{A}$ with a bilinear mapping $(a, b) \mapsto ab: \mathcal{A} \times \mathcal{A} \rightarrow \mathcal{A}$ and with a special nonzero element $1 \in \mathcal{A}$ such that

$$(ab)c = a(bc) \quad \text{and} \quad 1a = a = a1. \quad (1.1)$$

When we define the *multiplication* as the linear mapping $m: \mathcal{A} \otimes \mathcal{A} \rightarrow \mathcal{A}$ such that $m(a \otimes b) = ab$ and the *unit* as the linear mapping $\eta: \mathbb{C} \rightarrow \mathcal{A}$ such that $\eta(1) = 1_{\mathcal{A}}$, then we can rephrase (1.1) as

$$m \circ (m \otimes \text{id}) = m \circ (\text{id} \otimes m), \quad (1.2)$$

$$m \circ (\eta \otimes \text{id}) = \text{id} = m \circ (\text{id} \otimes \eta). \quad (1.3)$$

Here the two sides of (1.2) are linear mappings from $\mathcal{A} \otimes \mathcal{A} \otimes \mathcal{A}$ to $\mathcal{A}$, while the three parts of (1.3) are linear mappings from $\mathcal{A}$ to $\mathcal{A}$. In (1.3) we identify $\mathbb{C} \otimes \mathcal{A}$ with $\mathcal{A}$ by identifying $c \otimes a$ with ca ($c \in \mathbb{C}$, $a \in \mathcal{A}$). Thus $(\eta \otimes \text{id})(a) = (\eta \otimes \text{id})(1 \otimes a) = 1_{\mathcal{A}} \otimes a$.

Definition 1.1 A *coassociative coalgebra with counit* (or shortly a *coalgebra*) is a linear space $\mathcal{A}$ with linear mappings $\Delta: \mathcal{A} \rightarrow \mathcal{A} \otimes \mathcal{A}$ (*comultiplication*) and $\varepsilon: \mathcal{A} \rightarrow \mathbb{C}$ (*counit*) (nonzero) such that

$$(\Delta \otimes \text{id}) \circ \Delta = (\text{id} \otimes \Delta) \circ \Delta \quad (\text{coassociativity}), \quad (1.4)$$

$$(\varepsilon \otimes \text{id}) \circ \Delta = \text{id} = (\text{id} \otimes \varepsilon) \circ \Delta. \quad (1.5)$$

Here the two sides of (1.4) are linear mappings from $\mathcal{A}$ to $\mathcal{A} \otimes \mathcal{A} \otimes \mathcal{A}$, while the three parts of (1.5) are linear mappings from $\mathcal{A}$ to $\mathcal{A}$. Here we used again the identification of $\mathbb{C} \otimes \mathcal{A}$ or $\mathcal{A} \otimes \mathbb{C}$ with $\mathcal{A}$.

If $\mathcal{A}$ is an algebra with identity element 1 then $\mathcal{A} \otimes \mathcal{A}$ naturally becomes an algebra with identity element $1 \otimes 1$ if we define the product of $a \otimes b$ and $c \otimes d$ as $ac \otimes bd$.

When we speak of an algebra homomorphism then we will always mean that the homomorphism mapping also sends 1 to 1 (*unital* algebra homomorphism).

Definition 1.2 A *bialgebra* is a linear space $\mathcal{A}$ which has the structure of both an algebra and a coalgebra such that the mappings $\Delta: \mathcal{A} \rightarrow \mathcal{A} \otimes \mathcal{A}$ and $\varepsilon: \mathcal{A} \rightarrow \mathbb{C}$ are algebra homomorphisms.

Definition 1.3 A *Hopf algebra* is a bialgebra $\mathcal{A}$ together with a linear mapping $S: \mathcal{A} \rightarrow \mathcal{A}$ (*antipode*) such that

$$m \circ (S \otimes \text{id}) \circ \Delta = \eta \circ \varepsilon = m \circ (\text{id} \otimes S) \circ \Delta \quad (1.6)$$

(identities of linear mappings from $\mathcal{A}$ to $\mathcal{A}$), i.e.,

$$(m \circ (S \otimes \text{id}) \circ \Delta)(a) = \varepsilon(a) 1 = (m \circ (\text{id} \otimes S) \circ \Delta)(a), \quad a \in \mathcal{A}. \quad (1.7)$$

If V is a linear space then the *flip operator* $\sigma: V \otimes V \rightarrow V \otimes V$ will be the linear operator such that $\sigma(v_1 \otimes v_2) = v_2 \otimes v_1$ for $v_1, v_2 \in V$. Note that an algebra $\mathcal{A}$ is commutative iff $m \circ \sigma = m$. By analogy, we define a coalgebra $\mathcal{A}$ to be *cocommutative* if $\sigma \circ \Delta = \Delta$.

Example 1.4 Let G be a group and let $\mathcal{A} := \text{Fun}(G)$ be the algebra (under pointwise multiplication) of all complex-valued functions on G . To some extent, the algebra $\text{Fun}(G \times G)$ of all complex-valued functions on $G \times G$ can be viewed as the tensor product $\mathcal{A} \otimes \mathcal{A}$. Just write $(a \otimes b)(x, y) := a(x)b(y)$ if $a, b \in \mathcal{A}$ and $x, y \in G$. However, $\text{Fun}(G \times G)$ is not the algebraic tensor product of $\text{Fun}(G)$ with $\text{Fun}(G)$ except if G is a finite group. Let us for the moment not worry about this. Since $(m(a \otimes b))(x) = (ab)(x) = a(x)b(x) = (a \otimes b)(x, x)$ if $a, b \in \text{Fun}(G)$, we can write

$$(m(F))(x) = F(x, x), \quad F \in \text{Fun}(G \times G), \quad x \in G. \quad (1.8)$$

Now define the comultiplication, counit and antipode by

$$(\Delta(a))(x, y) := a(xy), \quad a \in \mathcal{A}, \quad x, y \in G, \quad (1.9)$$

$$\varepsilon(a) := a(e), \quad a \in \mathcal{A}, \quad (1.10)$$

$$(S(a))(x) := a(x^{-1}), \quad a \in \mathcal{A}, \quad x \in G. \quad (1.11)$$

The general philosophy here is that all properties of and information about the group can be stored in the algebra $\text{Fun}(G)$. Thus group multiplication, group identity and group inverse are described on the level of $\text{Fun}(G)$ by Δ , ε and S , respectively.

Evidently, Δ and ε are algebra homomorphisms and S is a linear mapping (in this example also an algebra homomorphism, while $S^2 = \text{id}$). Furthermore, the group axioms yield the Hopf algebra axioms (1.4), (1.5) and (1.6). This can be seen for the first identity in (1.7) by observing from (1.9) and (1.11) that $((S \otimes \text{id}) \circ \Delta)(a)(x, y) = a(x^{-1}y)$ and next from (1.8) that the left hand side of (1.7) evaluated in $x \in G$ equals $a(x^{-1}x) = a(e)$. By (1.10) the evaluation of the middle part of (1.7) in x yields the same.

Cocommutativity $\sigma \circ \Delta = \Delta$ would be equivalent here to $a(xy) = a(yx)$ for all $a \in \text{Fun}(G)$. Thus $\text{Fun}(G)$ is cocommutative iff the group G is abelian.

Everything above holds rigorously with algebraic tensor products if G is a finite group. Now suppose that G is a subgroup of $SL(n, \mathbb{C})$ (the group of complex $n \times n$ matrices of determinant 1). Usually (but not necessarily), this subgroup G may be thought to be *algebraic*, i.e. closed in the Zariski topology. Let $\text{Pol}(G)$ consist of all complex-valued functions depending on $x \in G$ which can be written as polynomials in the matrix elements x_{ij} of x . Let $\text{Pol}(G \times G)$ consist of all complex-valued functions of $(x, y) \in G \times G$ which can be written as polynomials in the matrix elements x_{ij} and y_{ij} of x and y . Then $\text{Pol}(G \times G)$ can be identified with $\text{Pol}(G) \otimes \text{Pol}(G)$. Now $\mathcal{A} := \text{Pol}(G)$ becomes a Hopf algebra in the algebraic sense with the above defined operations.

Let t_{ij} be the element of $\mathcal{A} = \text{Pol}(G)$ such that $t_{ij}(x) = x_{ij}$ ($x \in G$). Then the t_{ij} form a set of generators of the algebra $\mathcal{A}$. In fact, $\mathcal{A}$ is the quotient algebra obtained when the free abelian algebra with 1 generated by the t_{ij} is divided by the ideal of all elements in this free algebra which vanish on G . The Hopf algebra operations can now be specified by defining them for the generators. Thus, since

$$(\Delta(t_{ij}))(x, y) = t_{ij}(xy) = (xy)_{ij} = \sum_{k=1}^n x_{ik} y_{kj} = \sum_{k=1}^n t_{ik}(x) t_{kj}(y),$$

we have

$$\Delta(t_{ij}) = \sum_{k=1}^n t_{ik} \otimes t_{kj}, \quad (1.12)$$

and similarly

$$\varepsilon(t_{ij}) = \delta_{ij}, \quad S(t_{ij}) = T_{ji}, \quad (1.13)$$

where T_{ji} is the cofactor of the (ji) th entry in the matrix $(t_{kl})_{k,l=1,\dots,n}$.

The following notation is often useful. Let $\mathcal{A}$ be a coalgebra. If $a \in \mathcal{A}$ then we can choose sets of elements $a_{(1)i}$ and $a_{(2)i}$ in $\mathcal{A}$ (i running over a finite set) such that $\Delta(a) = \sum_i a_{(1)i} \otimes a_{(2)i}$. We write this symbolically as

$$\Delta(a) = \sum_{(a)} a_{(1)} \otimes a_{(2)}, \quad a \in \mathcal{A}. \quad (1.14)$$

Similarly, we write

$$(\Delta \otimes \text{id})(\Delta(a)) = \sum_{(a)} a_{(1)} \otimes a_{(2)} \otimes a_{(3)}, \quad a \in \mathcal{A}.$$

This notation is justified by the coassociativity (1.4).

Now we can express identities for Hopf algebras which involve comultiplication by this notation. For instance, (1.5) (applied to $a \in \mathcal{A}$) and (1.7) can be written as

$$\sum_{(a)} \varepsilon(a_{(1)}) a_{(2)} = a = \sum_{(a)} \varepsilon(a_{(2)}) a_{(1)}, \quad (1.15)$$

$$\sum_{(a)} S(a_{(1)}) a_{(2)} = \varepsilon(a) 1 = \sum_{(a)} a_{(1)} S(a_{(2)}). \quad (1.16)$$

Proposition 1.5 Let $\mathcal{A}$ be a bialgebra. If an antipode S exists such that $\mathcal{A}$ becomes a Hopf algebra then S is unique.

Proof Define a *convolution product* $F * G$ of linear mappings F and G from the bialgebra $\mathcal{A}$ to itself by

$$(F * G)(a) := (m \circ (F \otimes G) \circ \Delta)(a) = \sum_{(a)} F(a_{(1)}) G(a_{(2)}), \quad a \in \mathcal{A}. \quad (1.17)$$

Coassociativity of Δ and associativity of m then show that, under this convolution operation, $\text{End}(\mathcal{A})$ is an associative algebra. The properties of counit and unit yield $\eta \circ \varepsilon$ as an identity element for this algebra. The antipode property (1.6) can now be interpreted as

$$S * \text{id} = \eta \circ \varepsilon = \text{id} * S. \quad (1.18)$$

Thus, if S exists then it is the two-sided inverse of id in this convolution algebra, and therefore unique. $\square$

Because of (1.18) we can write the antipode as $S = S * \text{id} * S$. This can be used in order to derive further properties of S . In particular, the next proposition states that S is unital, counital, anti-multiplicative and anti-comultiplicative.

Proposition 1.6 Let $\mathcal{A}$ be a Hopf algebra. Then, for $a, b \in \mathcal{A}$,

$$S(1) = 1, \quad (1.19)$$

$$\varepsilon(S(a)) = \varepsilon(a), \quad (1.20)$$

$$S(ab) = S(b) S(a), \quad (1.21)$$

$$\Delta(S(a)) = ((S \otimes S) \circ \sigma \circ \Delta)(a) = \sum_{(a)} S(a_{(2)}) \otimes S(a_{(1)}). \quad (1.22)$$

Proof Formula (1.19) follows by putting $a := 1$ in (1.7), while (1.20) follows from (1.15) and (1.16). For the proof of (1.21) write

$$\begin{aligned} S(b) S(a) &= \sum_{(a),(b)} S(b_{(1)}) S(a_{(1)}) (\varepsilon(a_{(2)} b_{(2)}) 1) \\ &= \sum_{(a),(b)} S(b_{(1)}) S(a_{(1)}) a_{(2)} b_{(2)} S(a_{(3)} b_{(3)}) \\ &= \sum_{(a),(b)} \varepsilon(a_{(1)}) \varepsilon(b_{(1)}) S(a_{(2)} b_{(2)}) = S(ab). \end{aligned}$$

In the first identity (1.15) was applied twice. In the second identity we used (1.16) with $a := a_{(2)} b_{(2)}$ and also the fact that Δ is an algebra homomorphism. In the third identity we used (1.16) twice. The fourth identity contains two further applications of (1.15).

The proof of (1.22) is given by

$$\begin{aligned} \sum_{(a)} S(a_{(2)}) \otimes S(a_{(1)}) &= \sum_{(a)} \varepsilon(a_{(3)}) S(a_{(2)}) \otimes S(a_{(1)}) \\ &= \sum_{(a)} (S(a_{(2)}) \otimes S(a_{(1)})) (\varepsilon(a_{(3)}) 1 \otimes 1) \\ &= \sum_{(a)} (S(a_{(2)}) \otimes S(a_{(1)})) (a_{(3)} \otimes a_{(4)}) \Delta(S(a_{(5)})) \\ &= \sum_{(a)} (S(a_{(2)}) a_{(3)} \otimes S(a_{(1)}) a_{(4)}) \Delta(S(a_{(5)})) \\ &= \sum_{(a)} \varepsilon(a_{(2)}) (1 \otimes S(a_{(1)}) a_{(3)}) \Delta(S(a_{(4)})) \\ &= \sum_{(a)} (1 \otimes S(a_{(1)}) a_{(2)}) \Delta(S(a_{(3)})) \\ &= \sum_{(a)} \varepsilon(a_{(1)}) \Delta(S(a_{(2)})) = \Delta(S(a)). \end{aligned}$$

For the third identity above, apply Δ to the second identity in (1.16).

Example 1.7 Let $\mathfrak{g}$ be a complex Lie algebra and $\mathcal{A} := \mathcal{U}(\mathfrak{g})$ its universal enveloping algebra. $\mathcal{U}(\mathfrak{g})$ is defined as the quotient algebra $\mathcal{T}(\mathfrak{g})/J$, where $\mathcal{T}(\mathfrak{g})$ is the tensor algebra of $\mathfrak{g}$ and J is the ideal in $\mathcal{T}(\mathfrak{g})$ which is generated by the elements $XY - YX - [X, Y]$ ($X, Y \in \mathfrak{g}$). We define Δ , ε and S first on $\mathfrak{g}$:

$$\Delta(X) := X \otimes 1 + 1 \otimes X, \quad \varepsilon(X) := 0, \quad S(X) := -X \quad \text{for } X \in \mathfrak{g}. \quad (1.23)$$

On $\mathbb{C}$ we declare the operators to be unital. Now we can check the coalgebra axioms and the antipode axiom already for the operators acting on $\mathbb{C} \oplus \mathfrak{g}$. In fact, the counit and antipode axiom force ε and S to be as in (1.23). Next we extend Δ and ε to $\mathcal{T}(\mathfrak{g})$ as algebra homomorphisms and S as anti-algebra homomorphism. Now check that $\Delta(J) \subset \mathcal{T}(\mathfrak{g}) \otimes J + J \otimes \mathcal{T}(\mathfrak{g})$, $\varepsilon(J) = 0$ and $S(J) \subset J$. This allows us to consider Δ , ε and S as operators on $\mathcal{U}(\mathfrak{g})$. Finally we have to check the Hopf algebra axioms on all of $\mathcal{U}(\mathfrak{g})$, by using that they are already satisfied on a subspace of generators. Note that $\mathcal{A}$ is cocommutative, but generally not commutative, unless $\mathfrak{g}$ is an abelian Lie algebra. Note that the antipode satisfies $S^2 = \text{id}$.

Remark 1.8 A further motivation for the concept of comultiplication is the wish to construct tensor products of representations of algebras. Suppose $\mathcal{A}$ is an algebra. Let π_1 and π_2 be algebra representations of $\mathcal{A}$ on finite dimensional linear spaces V_1 and V_2 resp., i.e., algebra homomorphisms of $\mathcal{A}$ to the algebras of linear endomorphisms of V_1 and V_2 respectively. Then $a_1 \otimes a_2 \mapsto \pi_1(a_1) \otimes \pi_2(a_2): \mathcal{A} \otimes \mathcal{A} \rightarrow \text{End}(V_1 \otimes V_2)$ is a representation of $\mathcal{A} \otimes \mathcal{A}$ on $V_1 \otimes V_2$. In order to obtain from this representation a representation of $\mathcal{A}$ on $V_1 \otimes V_2$, we need an algebra homomorphism from $\mathcal{A}$ to $\mathcal{A} \otimes \mathcal{A}$. Without further structure on $\mathcal{A}$ there is no canonical method for this. However, if $\mathcal{A}$ is a Hopf algebra, the desired mapping is provided by the comultiplication Δ . Then we can define the tensor product representation $\pi_1 \otimes \pi_2$ of $\mathcal{A}$ as the composition $(\pi_1 \otimes \pi_2) \circ \Delta$. Associativity of this tensor product is precisely assured by the coassociativity (1.4) of Δ .

Other Hopf algebra axioms are also meaningful in this context. For instance, the counit ε gives a one-dimensional algebra representation of $\mathcal{A}$. Then the counit axiom (1.5) implies that, for each finite dimensional algebra representation π of $\mathcal{A}$, we have that $\pi \otimes \varepsilon = \pi = \varepsilon \otimes \pi$. Also, if π is an algebra representation of $\mathcal{A}$ on a finite dimensional linear space V and if V^* is the linear dual of V , then we can define an algebra representation π^* of $\mathcal{A}$ on V^* by

$$\langle \pi^*(a) v^*, v \rangle := \langle v^*, \pi(S(a)) v \rangle, \quad v \in V, \quad v^* \in V^*.$$

Recall that the tensor product of two representations π_1 and π_2 of a Lie algebra $\mathfrak{g}$ is provided by $((\pi_1 \otimes \pi_2)(X))(v_1 \otimes v_2) = (\pi_1(X) v_1) \otimes v_2 + v_1 \otimes (\pi_2(X) v_2)$. Compare this with the comultiplication for $\mathfrak{g}$ as given by (1.23).

One should be aware that the antipode in a Hopf algebra $\mathcal{A}$ is not necessarily invertible as a linear mapping from $\mathcal{A}$ to itself.

We may consider $\mathcal{A}$ also as a Hopf algebra with opposite multiplication $m \circ \sigma$ and opposite comultiplication $\sigma \circ \Delta$, and ε , η and S unchanged. Then S is a Hopf algebra homomorphism from $\mathcal{A}$ with the old Hopf algebra structure to $\mathcal{A}$ with the new one. However, if we change only one of the two operations of multiplication and comultiplication into its opposite then we still have a bialgebra, but this will be a Hopf algebra if and only if S is invertible. The new antipode will then be precisely

S^{-1} . Indeed, let $\mathcal{A}'$ be $\mathcal{A}$ equipped with opposite multiplication and suppose $\mathcal{A}'$ is a Hopf algebra with antipode S' . Now apply S to

$$\sum_{(a)} a_{(2)} S'(a_{(1)}) = \varepsilon(a)1 = \sum_{(a)} S'(a_{(2)}) a_{(1)}.$$

This yields that SS' is the two-sided inverse of S under convolution. Hence $SS' = \text{id}$, and also $S'S = \text{id}$ by reverting the roles of S and S' . Conversely, if S is invertible, then define $S' := S^{-1}$ and show that S' is an antipode for $\mathcal{A}'$.

Recall that an (associative) algebra $\mathcal{A}$ (with 1) is a $*$ -algebra if there is a mapping $a \mapsto a^*: \mathcal{A} \rightarrow \mathcal{A}$ (an *involution*) such that $(a^*)^* = a$, $(a+b)^* = a^* + b^*$, $(\lambda a)^* = \bar{\lambda} a^*$, $(ab)^* = b^* a^*$, $1^* = 1$. If $\mathcal{A}$ is a $*$ -algebra then $\mathcal{A} \otimes \mathcal{A}$ becomes a $*$ -algebra with $(a \otimes b)^* := a^* \otimes b^*$. If $\mathcal{A}$ and $\mathcal{B}$ are $*$ -algebras then $F: \mathcal{A} \rightarrow \mathcal{B}$ is called a $*$ -homomorphism if F is an algebra homomorphism such that $F(a^*) = (F(a))^*$.

Definition 1.9 A Hopf $*$ -algebra is a Hopf algebra $\mathcal{A}$ which, as an algebra, is also a $*$ -algebra such that $\Delta: \mathcal{A} \rightarrow \mathcal{A} \otimes \mathcal{A}$ and $\varepsilon: \mathcal{A} \rightarrow \mathbb{C}$ are $*$ -homomorphisms.

Originally, the property stated in the next Proposition was part of the definition of a Hopf $*$ -algebra. I thank S. Zakrzewski for helpful correspondence about this.

Proposition 1.10 In a Hopf $*$ -algebra $\mathcal{A}$ we have

$$S \circ * \circ S \circ * = \text{id}. \quad (1.24)$$

In particular, S is invertible with inverse $* \circ S \circ *$.

Proof Apply (1.6) to a^* and next apply the involution to all members of (1.6). This shows that $* \circ S \circ *$ is an antipode for $\mathcal{A}'$, i.e., for $\mathcal{A}$ with opposite multiplication. Hence $* \circ S \circ * = S^{-1}$. $\square$

Example 1.11 Let G be a group and let $\mathcal{A} := \text{Fun}(G)$ as in Example 1.4 For $f \in \text{Fun}(G)$ define $f^*(x) := \overline{f(x)}$ ($x \in G$). Similarly, for $F \in \text{Fun}(G \times G)$ define $F^*(x, y) := \overline{F(x, y)}$ ($x, y \in G$). Then $\text{Fun}(G)$ and $\text{Fun}(G \times G)$ are commutative $*$ -algebras and $\Delta: \text{Fun}(G) \rightarrow \text{Fun}(G \times G)$ and $\varepsilon: \text{Fun}(G) \rightarrow \mathbb{C}$ are $*$ -homomorphisms. So, apart from the fact that Δ does not map $\text{Fun}(G)$ into $\text{Fun}(G) \otimes \text{Fun}(G)$, we have $\text{Fun}(G)$ as an example of a Hopf $*$ -algebra.

Example 1.12 Let G be a complex Lie group given as a closed connected subgroup of $SL(n, \mathbb{C})$ and let G_0 be a real connected Lie group and a real form of G . Then every $a \in \text{Pol}(G)$ is completely determined by its restriction to $\underline{G_0}$. Suppose that, for each $a \in \text{Pol}(G)$, there exists $a^* \in \text{Pol}(G)$ such that $a^*(x) = \overline{a(x)}$ for $x \in G_0$. Then the Hopf algebra $\text{Pol}(G)$ becomes a Hopf $*$ -algebra with this mapping $*$. Conversely,

if the Hopf algebra $\text{Pol}(G)$ is a Hopf $*$ -algebra then we may define a real form G_0 of G by $G_0 := \{x \in G \mid \overline{a(x)} = a^*(x) \quad \forall a \in \text{Pol}(G)\}$.

Thus, on the level of polynomial function algebras, a real Lie group is described by the algebra of functions on its complexification together with an involution for this algebra.

In particular, let the t_{ij} (cf. Example 1.4) be the generators of the algebra $\mathcal{A} = \text{Pol}(G)$. If we suppose that $G_0 \subset SL(n, \mathbb{R})$ then $t_{ij}^*(x) = \overline{t_{ij}(x)} = t_{ij}(x)$ for $x \in G_0$. Hence $t_{ij}^* = t_{ij}$. This gives the action of $*$ on the generators and next, by anti-linear homomorphic continuation, on all of $\mathcal{A}$. However, if $G_0 \subset SU(n)$ then $t_{ij}^*(x) = \overline{t_{ij}(x)} = t_{ji}(x^{-1}) = (S(t_{ji}))(x)$ if $x \in G_0$. Hence $t_{ij}^* = S(t_{ji})$. Again, the involution on all of $\mathcal{A}$ follows by anti-linear homomorphic continuation.

1.2. Duality for Hopf algebras. Let $\mathcal{A}$ be a Hopf algebra and let $\mathcal{A}^*$ be its algebraic linear dual, i.e., the space of all linear mappings $f: \mathcal{A} \rightarrow \mathbb{C}$. We will write $\langle f, a \rangle := f(a)$ ($f \in \mathcal{A}^*$, $a \in \mathcal{A}$). The algebraic tensor product $\mathcal{A}^* \otimes \mathcal{A}^*$ is a subspace of $(\mathcal{A} \otimes \mathcal{A})^*$ by the rule

$$\langle f \otimes g, a \otimes b \rangle = \langle f, a \rangle \langle g, b \rangle, \quad f, g \in \mathcal{A}^*, \quad a, b \in \mathcal{A}.$$

Unless $\mathcal{A}$ is finite dimensional, this will be a proper subspace. By duality, the Hopf algebra operations on $\mathcal{A}$ can be transferred to $\mathcal{A}^*$. We define

$$\langle fg, a \rangle := \langle f \otimes g, \Delta(a) \rangle = \sum_{(a)} \langle f, a_{(1)} \rangle \langle g, a_{(2)} \rangle, \quad (1.25)$$

$$\langle \Delta(f), a \otimes b \rangle := \langle f, ab \rangle, \quad (1.26)$$

$$\langle 1_{\mathcal{A}^*}, a \rangle := \varepsilon_{\mathcal{A}}(a), \quad (1.27)$$

$$\varepsilon_{\mathcal{A}^*}(f) := \langle f, 1_{\mathcal{A}} \rangle, \quad (1.28)$$

$$\langle S(f), a \rangle := \langle f, S(a) \rangle. \quad (1.29)$$

However, $\Delta(f)$, as defined by (1.26), will be an element of $(\mathcal{A} \otimes \mathcal{A})^*$ and not necessarily of $\mathcal{A}^* \otimes \mathcal{A}^*$. Still, with a suitable adaptation of the definition of Hopf algebra, the Hopf algebra axioms for $\mathcal{A}^*$ with the above operations can be verified in a straightforward way.

If $\mathcal{A}$ is moreover a Hopf $*$ -algebra then we can define an involution on $\mathcal{A}^*$ by

$$f^*(a) := \overline{\langle f, (S(a))^* \rangle}. \quad (1.30)$$

Note that (1.24) ensures that $(f^*)^* = f$. The antipode is needed in (1.30) in order to ensure that $(fg)^* = g^*f^*$. (The more simple definition $f^*(a) := \overline{\langle f, a^* \rangle}$ would make the involution on $\mathcal{A}^*$ multiplicative rather than anti-multiplicative.) The reader should verify that $\mathcal{A}^*$ thus indeed becomes a Hopf $*$ -algebra, with suitably adapted definition because of the fact that Δ does not necessarily map into $\mathcal{A}^* \otimes \mathcal{A}^*$.

Let $\mathcal{A}$ be a Hopf algebra. We define left and right algebra actions of $\mathcal{A}^*$ on $\mathcal{A}$:

$$f.a := (\text{id} \otimes f)(\Delta(a)) = \sum_{(a)} f(a_{(2)}) a_{(1)}, \quad (1.31)$$

$$a.f := (f \otimes \text{id})(\Delta(a)) = \sum_{(a)} f(a_{(1)}) a_{(2)}. \quad (1.32)$$

It is indeed an easy exercise to verify that

$$(fg).a = f.(g.a), \quad a.(fg) = (a.f).g, \quad 1_{\mathcal{A}^*}.a = a.1_{\mathcal{A}^*} = a.$$

An important property is the following. If $f \in \mathcal{A}^*$ such that $\Delta(f) \in \mathcal{A}^* \otimes \mathcal{A}^*$ and if we write $\Delta(f) = \sum_{(f)} f_{(1)} \otimes f_{(2)}$ then

$$f.(ab) = \sum_{(f)} (f_{(1)}.a) (f_{(2)}.b), \quad a, b \in \mathcal{A}, \quad (1.33)$$

$$(ab).f = \sum_{(f)} (a.f_{(1)}) (b.f_{(2)}), \quad a, b \in \mathcal{A}. \quad (1.34)$$

We give the proof of (1.33); the proof of (1.34) is similar.

$$\begin{aligned} f.(ab) &= \sum_{(a),(b)} a_{(1)} b_{(1)} f(a_{(2)} b_{(2)}) = \sum_{(a),(b),(f)} a_{(1)} f_{(1)}(a_{(2)}) b_{(1)} f_{(2)}(b_{(2)}) \\ &= \sum_{(f)} (f_{(1)}.a) (f_{(2)}.b). \end{aligned}$$

Definition 1.13 Two Hopf algebras $\mathcal{U}$ and $\mathcal{A}$ are said to be *Hopf algebras in duality* if there is a bilinear mapping $(u, a) \mapsto \langle u, a \rangle : \mathcal{U} \times \mathcal{A} \rightarrow \mathbb{C}$ such that (1.25)–(1.29) are satisfied when $\mathcal{U}$ is read instead of $\mathcal{A}^*$.

If $\mathcal{U}$ and $\mathcal{A}$ are moreover Hopf $*$ -algebras and if (1.30) also holds then we speak about *Hopf $*$ -algebras in duality*.

Two Hopf $(*)$ -algebras in duality $\mathcal{A}$ and $\mathcal{U}$ are said to be *Hopf $(*)$ -algebras in nondegenerate duality* (in fact in doubly nondegenerate duality) if the two following implications moreover hold: (i) $(\forall a \in \mathcal{A} \langle u, a \rangle = 0) \implies u = 0$, and (ii) $(\forall u \in \mathcal{U} \langle u, a \rangle = 0) \implies a = 0$.

If $\mathcal{U}$ and $\mathcal{A}$ are Hopf algebras in duality (not necessarily nondegenerate) then left and right actions of $\mathcal{U}$ on $\mathcal{A}$ as in (1.31) and (1.32) can still be defined.

If $\mathcal{A}$ is a Hopf $(*)$ -algebra and if $\mathcal{U}$ is a Hopf $(*)$ -subalgebra of $\mathcal{A}^*$, i.e., if $\mathcal{U}$ is a unital subalgebra of $\mathcal{A}^*$ such that $\Delta(\mathcal{U}) \subset \mathcal{U} \otimes \mathcal{U}$, $S(\mathcal{U}) \subset \mathcal{U}$ and, in case of a Hopf $*$ -subalgebra, $\mathcal{U}^* \subset \mathcal{U}$, then $\mathcal{U}$ and $\mathcal{A}$ are obviously Hopf $(*)$ -algebras in duality, while the duality is already nondegenerate on one side: if $u \in \mathcal{U}$ and if $\langle u, a \rangle = 0$ for all $a \in \mathcal{A}$ then $u = 0$.

If $\mathcal{U}$ and $\mathcal{A}$ are Hopf $(*)$ -algebras in nondegenerate duality then $\mathcal{U}$ can be viewed as a Hopf $(*)$ -subalgebra of $\mathcal{A}^*$ and $\mathcal{A}$ as a Hopf $(*)$ -subalgebra of $\mathcal{U}^*$. So we can write $\langle u, a \rangle = u(a) = a(u)$ for $u \in \mathcal{U}$ and $a \in \mathcal{A}$.

Example 1.14 Let G be a complex Lie group given as a closed connected subgroup of $SL(n, \mathbb{C})$ and let $\mathfrak{g}$ be its (complex) Lie algebra and $\mathcal{U} := \mathcal{U}(\mathfrak{g})$ the universal enveloping algebra of $\mathfrak{g}$. Then $\mathcal{A} := \text{Pol}(G)$ and $\mathcal{U}$ are Hopf algebras (cf. Examples 1.4 and 1.7). These Hopf algebras are naturally in duality with each other. For $X \in \mathfrak{g}$ and $a \in \text{Pol}(G)$ the pairing is given by

$$\langle X, a \rangle = \left. \frac{d}{dt} \right|_{t=0} a(\exp(tX)). \quad (1.35)$$

Then, for $f := X \in \mathfrak{g}$, equations (1.26)–(1.29) are satisfied in view of (1.35) and (1.23).

For the left and right actions of $X \in \mathfrak{g}$ on $a \in \text{Pol}(G)$ (cf. (1.31) and (1.32)) we then obtain

$$(X.a)(x) = \left. \frac{d}{dt} \right|_{t=0} a(x \exp(tX)), \quad (a.X)(x) = \left. \frac{d}{dt} \right|_{t=0} a(\exp(tX) x).$$

If $X := X_1 X_2 \dots X_k \in \mathcal{U}$, with $X_1, \dots, X_k \in \mathfrak{g}$, then

$$\begin{aligned} (X.a)(x) &= \left. \frac{\partial^k}{\partial t_1 \dots \partial t_k} \right|_{t_1, \dots, t_k=0} a(x \exp(t_1 X_1) \dots \exp(t_k X_k)), \\ (a.X)(x) &= \left. \frac{\partial^k}{\partial t_1 \dots \partial t_k} \right|_{t_1, \dots, t_k=0} a(\exp(t_1 X_1) \dots \exp(t_k X_k) x). \end{aligned}$$

So we get the familiar left or right action of $\mathcal{U}(\mathfrak{g})$ on smooth functions on G by left or right invariant differential operators. In particular, either of the two last formulas, when evaluated at $x := e$, yields the pairing $\langle X_1 \dots X_k, a \rangle$. The formula thus obtained is compatible with (1.35) and (1.25). The double non-degeneracy of the pairing can be easily verified.

Example 1.15 Let G , $\mathfrak{g}$, $\mathcal{A}$ and $\mathcal{U}$ be as in the previous example. Let the connected real Lie group G_0 be a real form of G and let $*$ be the corresponding involution on $\mathcal{A}$, by which $\mathcal{A}$ becomes a Hopf $*$ -algebra (cf. Example 1.12). Let $\mathfrak{g}_0$ be the (real) Lie algebra of G_0 . It is a real form of the complex Lie algebra $\mathfrak{g}$. Let us compute X^* (as defined by (1.30)) for $X \in \mathfrak{g}_0$ and let us see if $X^* \in \mathcal{U}$. For $a \in \mathcal{A}$ we have

$$\langle X^*, a \rangle = \overline{\langle X, (S(a))^* \rangle} = \overline{\left. \frac{d}{dt} \right|_{t=0} \overline{a(\exp(-tX))}} = -\langle X, a \rangle.$$

Here we used that $a^*(x) = \overline{a(x)}$ if $x \in G_0$ (cf. Example 1.12). Thus $X^* = -X$ for $X \in \mathfrak{g}_0$. In particular, $*$ maps the Lie algebra $\mathfrak{g}_0$ to itself. It follows that $*$ maps $\mathcal{U}$ to itself, so $\mathcal{U}$ is a Hopf $*$ -algebra in duality with $\mathcal{A}$.

Note that in the case $G_0 \subset SU(n)$ we have $\mathfrak{g}_0 \subset su(n)$, the Lie algebra of skew-hermitian matrices. So then the $*$ on $\mathfrak{g}$ induced by the $*$ -structure on $\mathcal{A}$ has the same effect as taking the adjoint of the matrix $X \in \mathfrak{g}$. But this is not necessarily true if $G_0 \subset SL(n, \mathbb{R})$.

1.3. Corepresentations.

Definition 1.16 Let $\mathcal{A}$ be a coalgebra. A *corepresentation* of $\mathcal{A}$ on a complex vector space V is defined as a linear mapping $t: V \rightarrow V \otimes \mathcal{A}$ such that

$$(t \otimes \text{id}) \circ t = (\text{id} \otimes \Delta) \circ t, \quad (\text{id} \otimes \varepsilon) \circ t = \text{id}. \quad (1.36)$$

Suppose that V is a finite dimensional vector space with basis $e_1, \dots, e_n$. Then t determines elements t_{ij} of $\mathcal{A}$ such that

$$t(e_j) = \sum_i e_i \otimes t_{ij}. \quad (1.37)$$

Then

$$\sum_i e_i \otimes \Delta(t_{ik}) = (\text{id} \otimes \Delta)(t(e_k)) = (t \otimes \text{id})(t(e_k)) = \sum_{j,k} e_i \otimes t_{ij} \otimes t_{jk}$$

and

$$\sum_i \varepsilon(t_{ij}) e_i = (\text{id} \otimes \varepsilon)(t(e_j)) = e_j.$$

Hence

$$\Delta(t_{ij}) = \sum_k t_{ik} \otimes t_{kj}, \quad \varepsilon(t_{ij}) = \delta_{ij}. \quad (1.38)$$

Conversely, if elements t_{ij} of $\mathcal{A}$ satisfy (1.38) and if t is defined by (1.37) on the vector space having the e_i as a basis then t is a corepresentation of $\mathcal{A}$ according to Definition 1.16. We call a matrix $(t_{ij})_{i,j=1,\dots,n}$ satisfying (1.38) a *matrix corepresentation* of $\mathcal{A}$.

Example 1.17 Let G be a group and π a representation of G on some vector space V . Let $\text{Fun}(G)$ be as in Example 1.4. To some extent, we can identify $V \otimes \text{Fun}(G)$ with $\text{Fun}(G; V)$ (the space of V -valued functions on G). Just write $(v \otimes a)(x) := a(x)v$ if $v \in V$, $a \in \mathcal{A}$, $x \in G$. This identification holds in the sense of algebraic tensor products if V is finite dimensional. Similarly, identify $V \otimes \text{Fun}(G) \otimes \text{Fun}(G)$ with $\text{Fun}(G \times G; V)$ by $(v \otimes a \otimes b)(x, y) := a(x)b(y)v$. Define $t(v) \in V \otimes \text{Fun}(G)$ ($v \in V$) as the V -valued function given by

$$(t(v))(x) := \pi(x)v, \quad x \in G.$$

Then t is a corepresentation of $\text{Fun}(G)$ on V . For this purpose, check equations (1.36). For instance, for the proof of the first identity of (1.36) we can show that

$$(((t \otimes \text{id}) \circ t)(v))(x, y) = \pi(x)\pi(y)v \quad \text{and} \quad (((\text{id} \otimes \Delta) \circ t)(v))(x, y) = \pi(xy)v.$$

Let us show the first of these two identities. Fix v . Then there are $v_j \in V$ and $a_j \in \text{Fun}(G)$ such that $(t(v))(x) = \pi(x)v = \sum_j a_j(x) v_j$, so $t(v) = \sum_j v_j \otimes a_j$. Then

$$((t \otimes \text{id}) \circ t)(v) = \sum_j (t \otimes \text{id})(v_j \otimes a_j) = \sum_j t(v_j) \otimes a_j.$$

Hence $((t \otimes \text{id}) \circ t)(v)(x, y)$

$$= \sum_j (t(v_j) \otimes a_j)(x, y) = \sum_j a_j(y) (t(v_j))(x) = \sum_j a_j(y) \pi(x) v_j = \pi(x) \pi(y) v.$$

Example 1.18 If t is a matrix representation of a group G then the matrix elements t_{ij} can be viewed as elements of $\text{Fun}(G)$. They satisfy

$$t_{ij}(xy) = \sum_k t_{ik}(x) t_{kj}(y), \quad t_{ij}(e) = \delta_{ij}.$$

Hence, by (1.38), t is a matrix corepresentation of $\text{Fun}(G)$. If t is a matrix representation of a group $G \subset SL(n, \mathbb{C})$ and if the t_{ij} are in $\text{Pol}(G)$ then t is a matrix corepresentation of $\text{Pol}(G)$. This holds in particular for the natural matrix representation of G defined by $t_{ij}(x) := x_{ij}$ ($x \in G$), cf. the end of Example 1.4.

If t is a matrix corepresentation of a Hopf algebra $\mathcal{A}$ then combination of (1.38) and (1.6) shows that

$$\sum_k S(t_{ik}) t_{kj} = \delta_{ij} 1 = \sum_k t_{ik} S(t_{kj}). \quad (1.39)$$

For a corepresentation t of $\mathcal{A}$ on V we may use a symbolic notation analogous to (1.14):

$$t(v) = \sum_{(v)} v_{(1)} \otimes v_{(2)}, \quad v \in V, \quad (1.40)$$

which means that we have a sum of elements $t_{(1)i} \otimes t_{(2)i}$, where $t_{(1)i} \in V$ and $t_{(2)i} \in \mathcal{A}$. Then (1.36) allows us to combine (1.40) with (1.14) and to write

$$((t \otimes \text{id}) \circ t)(v) = \sum_{(v)} v_{(1)} \otimes v_{(2)} \otimes v_{(3)} = ((\text{id} \otimes \Delta) \circ t)(v).$$

Definition 1.19 Let t be a corepresentation of a Hopf $*$ -algebra $\mathcal{A}$ on a vector space V with hermitian inner product $(\cdot, \cdot)$. We call the corepresentation t *unitary* if

$$\sum_{(v), (w)} (v_{(1)}, w_{(1)}) w_{(2)}^* v_{(2)} = (v, w) 1, \quad v, w \in V. \quad (1.41)$$

Condition (1.41) can be equivalently stated as

$$\sum_{(v)} (v_{(1)}, w) S(v_{(2)}) = \sum_{(w)} (v, w_{(1)}) w_{(2)}^*, \quad v, w \in V. \quad (1.42)$$

We prove the implication (1.41) $\Rightarrow$ (1.42), and leave the other direction to the reader. Indeed,

$$\begin{aligned} \sum_{(v)} (v_{(1)}, w) S(v_{(2)}) &= \sum_{(v), (w)} (v_{(1)}, w_{(1)}) w_{(2)}^* v_{(2)} S(v_{(3)}) \\ &= \sum_{(v), (w)} (v_{(1)}, w_{(1)}) w_{(2)}^* \varepsilon(v_{(2)}) = \sum_{(w)} (v, w_{(1)}) w_{(2)}^*. \end{aligned}$$

If V is finite dimensional with an orthonormal basis $e_1, \dots, e_n$ then (1.41) implies for the corresponding matrix corepresentation (t_{ij}) that

$$\sum_i t_{il}^* t_{ik} = \sum_{i,j} (e_i, e_j) t_{jl}^* t_{ik} = (e_k, e_l) 1 = \delta_{kl} 1.$$

In combination with (1.39) we readily obtain:

Proposition 1.20 Let (t_{ij}) be a matrix corepresentation of a Hopf $*$ -algebra $\mathcal{A}$ and let t be the corresponding corepresentation of $\mathcal{A}$ on the inner product space which has $e_1, \dots, e_n$ as an orthonormal basis. Then the following conditions are equivalent.

- (a) t is a unitary corepresentation.
- (b) $\sum_k t_{ki}^* t_{kj} = \delta_{ij} 1$.
- (c) $S(t_{ij}) = t_{ji}^*$.
- (d) $\sum_k t_{ik} t_{jk}^* = \delta_{ij} 1$.

In Example 1.18 we viewed a matrix representation t of a group G as a matrix corepresentation of $\text{Fun}(G)$. If we have a $*$ -operation on $\text{Fun}(G)$ as in Example 1.11, then it follows from Proposition 1.20 that t is unitary as a matrix representation of G iff it is unitary as a matrix corepresentation of $\text{Fun}(G)$.

If π is a unitary representation of G on an inner product space V and if t is the corepresentation of $\text{Fun}(G)$ associated with π according to Example 1.17 then equations (1.41) and (1.42) are rewritten versions of

$$(\pi(x)v, \pi(x)w) = (v, w) \quad \text{and} \quad (\pi(x^{-1})v, w) = (v, \pi(x)w), \quad x \in G_0.$$

Remark 1.21 If t is a corepresentation of $\mathcal{A}$ on a vector space V then t gives rise to an algebra representation π of $\mathcal{A}^*$ on V :

$$\pi(f)v := (\text{id} \otimes f)(t(v)) = \sum_{(v)} f(v_{(2)}) v_{(1)}. \quad (1.43)$$

Suppose V is finite dimensional with basis $e_1, \dots, e_n$ and let the corresponding matrix elements t_{ij} be given by (1.37). Combination with (1.43) gives that

$$\pi(f) e_j = \sum_i f(t_{ij}) e_i.$$

Hence π can then be written as a matrix representation $\pi: f \mapsto (\pi_{ij}(f))$ of $\mathcal{A}$ with $\pi_{ij}(f) := f(t_{ij})$.

If t is a unitary matrix corepresentation of a Hopf $*$ -algebra $\mathcal{A}$ and if π is the corresponding matrix representation of $\mathcal{A}^*$ then π is a $*$ -representation. Indeed,

$$\pi_{ij}(f^*) = f^*(t_{ij}) = \overline{f((S(t_{ij}))^*)} = \overline{f(t_{ji})} = \overline{\pi_{ji}(f)}.$$

Definition 1.22 Let t be a corepresentation of a coalgebra $\mathcal{A}$ on a vector space V .

- (a) A linear subspace W of V is called *invariant* if $t(W) \subset W \otimes \mathcal{A}$.
- (b) The corepresentation t is called *irreducible* if V and $\{0\}$ are the only invariant subspaces of V .
- (c) Let s be another corepresentation of $\mathcal{A}$ on a vector space W . A linear operator $L: V \rightarrow W$ is called an *intertwining operator* for t and s if

$$s \circ L = (L \otimes \text{id}) \circ t. \tag{1.44}$$

The corepresentations t and s are called *equivalent* if there exists a bijective intertwining operator for t and s .

Observe that the relation of equivalence between corepresentations is an equivalence relation. One easily checks that two matrix corepresentations (t_{ij}) and (s_{ij}) of a coalgebra $\mathcal{A}$ are equivalent iff there is an invertible square complex matrix L such that $Lt = sL$ (where the products are matrix products). Furthermore, the matrix corepresentation t is irreducible iff t is not equivalent to a matrix corepresentation of block form $\begin{pmatrix} * & * \\ 0 & * \end{pmatrix}$.

Proposition 1.23 Let t be a unitary corepresentation of a Hopf $*$ -algebra on a finite dimensional Hilbert space V .

- (a) Let W be an invariant subspace of V . Then the orthoplement of W in V is also invariant.
- (b) V is a direct sum of invariant subspaces on each of which the restriction of t is an irreducible corepresentation of $\mathcal{A}$.

Proof For the proof of (a) assume that v is orthogonal to W . Write $t(v) = \sum_{(v)} v_{(1)} \otimes v_{(2)}$ such that the $v_{(2)}$ are linearly independent. Apply (1.42) with $w \in W$. Then, by invariance of W , the right hand side of (1.42) will be 0. Hence the left hand side will be 0, so $\sum_{(v)} (v_{(1)}, w) v_{(2)} = 0$. Thus the $(v_{(1)}, w)$ will be zero. So the $v_{(1)}$ will be in the orthoplement of W . Part (b) follows by iteration of (a). $\square$

Lemma 1.24 Let $L: V \rightarrow W$ be an intertwining operator for corepresentations t and s of a coalgebra $\mathcal{A}$ on finite dimensional spaces V resp. W .

- (a) The image $L(V)$ and the null space $L^{-1}(0)$ are invariant subspaces of W and V , respectively.
- (b) (*first Schur lemma*) If t and s are irreducible then $L = 0$ or L is bijective.
- (c) (*second Schur lemma*) If $V = W$, $t = s$ and t is irreducible then $L = \lambda I$ for some complex λ .

Proof In (a) the invariance of $L(V)$ follows immediately from (1.44). In order to prove the invariance of $L^{-1}(0)$, let $Lv = 0$. Then, by (1.44) and (1.40),

$$0 = (L \otimes \text{id})(t(v)) = \sum_{(v)} L(v_{(1)}) \otimes v_{(2)}.$$

Now choose the $v_{(2)}$ linearly independent. Then the $L(v_{(1)})$ will be 0. Hence, the $v_{(1)}$ in (1.40) will belong to $L^{-1}(0)$. Parts (b) and (c) now follow easily, completely analogous to the proof of the classical Schur lemmas. $\square$

Lemma 1.25 Let $\mathcal{A}$ be a coalgebra. Let t be a corepresentation of $\mathcal{A}$ on a finite dimensional vector space V . Suppose that V is a direct sum of subspaces V_i ($i = 1, \dots, n$) and that each V_i is a direct sum of subspaces W_{ij} ($j = 1, \dots, m_i$) and that there are mutually inequivalent irreducible corepresentations $t_1, \dots, t_n$ such that each subspace W_{ij} is invariant and t restricted to W_{ij} is equivalent to t_i . Let U be a nonzero invariant subspace of V such that t restricted to U is an irreducible corepresentation s . Then, for some i , $U \subset V_i$ and s is equivalent to t_i .

Proof Let $P_{ij}: V \rightarrow W_{ij}$ be the projection operator which is identity on W_{ij} and 0 on the other W_{kl} . Let π_{ij} be the restriction of P_{ij} to U . Then $\pi_{ij}: U \rightarrow W_{ij}$ is an intertwining operator for s and t_i . Hence, by Lemma 1.24(b), it is either 0 or bijective. Hence, if π_{ij} and π_{kl} are nonzero, then t_i , s and t_k must be equivalent corepresentations, so $i = k$. Hence, there is an i such that

$$U \subset \bigoplus_{k,l} \pi_{kl}(U) = \bigoplus_{\substack{k,l \\ \pi_{kl} \neq 0}} W_{kl} \subset \bigoplus_{j=1}^{m_i} W_{ij} = V_i$$

and s is equivalent to t_i . $\square$

If $\mathcal{A}$ and $\mathcal{B}$ are coalgebras then $\mathcal{A} \otimes \mathcal{B}$ becomes a coalgebra with comultiplication given by $\Delta(a \otimes b) := \sum_{(a),(b)} (a_{(1)} \otimes b_{(1)}) \otimes (a_{(2)} \otimes b_{(2)})$.

Lemma 1.26 Let $\mathcal{A}$ and $\mathcal{B}$ be coalgebras. Let s and t be irreducible corepresentations of $\mathcal{A}$ resp. $\mathcal{B}$ on finite dimensional vector spaces V and W . Define the

corepresentation r of the coalgebra $\mathcal{A} \otimes \mathcal{B}$ on $V \otimes W$ as the tensor product of the corepresentations s and t :

$$r(v \otimes w) = \sum_{(v), (w)} (v_{(1)} \otimes w_{(1)}) \otimes (v_{(2)} \otimes w_{(2)}). \quad (1.45)$$

Then r is irreducible.

Proof Let U be a nonzero invariant irreducible subspace of $V \otimes W$ with respect to r . It will readily follow that $U = V \otimes W$ if we can show that U contains a nonzero element of the form $v \otimes w$. Define the corepresentation $\tilde{s}$ of $\mathcal{A}$ on $V \otimes W$ by

$$\tilde{s}(v \otimes w) := \sum_{(v)} (v_{(1)} \otimes w) \otimes v_{(2)}.$$

Now, because $\tilde{s} = (\text{id} \otimes \text{id} \otimes \text{id} \otimes \varepsilon_{\mathcal{B}}) \circ r$, we see that U is invariant with respect to $\tilde{s}$, so U contains an invariant subspace U_0 on which the restriction s_0 of $\tilde{s}$ is an irreducible corepresentation of $\mathcal{A}$. Choose a basis $f_1, \dots, f_n$ for W . Let $P_j: V \otimes W \rightarrow V$ be the operator which sends $v \otimes f_j$ to v and is 0 on the other $V \otimes f_k$. Let π_j be the restriction of P_j to U_0 . Then π_j is an intertwining operator for s_0 and s . By Lemma 1.24(b) π_j is bijective or 0. Now π_j is nonzero for some j ; after rearranging the f_j we may assume that $\pi_1 \neq 0$. Then, for each j , $\pi_j \circ \pi_1^{-1}: V \rightarrow V$ is an intertwining operator for s and s . Thus, by Lemma 1.24(c), $\pi_j \circ \pi_1^{-1} = \lambda_j I$ for some complex λ . We conclude that $u \in U_0$ can be written as $u = \sum_{j=1}^n \pi_j(u) \otimes f_j = \pi_1(u) \otimes (\sum_{j=1}^n \lambda_j f_j)$. $\square$

Definition 1.27 Let $(t_{ij})_{i,j=1,\dots,n}$ be a matrix corepresentation of a Hopf algebra $\mathcal{A}$. Then, because of (1.20) and (1.22), $(S(t_{ji}))_{i,j=1,\dots,n}$ is also a matrix corepresentation of $\mathcal{A}$. We denote it by (t'_{ij}) and we call it the *contragredient corepresentation* of t .

Note that, if the matrix corepresentations t and s are equivalent with intertwining operator given by the matrix A then t' and s' are equivalent with intertwining operator given by the matrix ${}^t A^{-1}$. If S is invertible (for instance if $\mathcal{A}$ is a Hopf $*$ -algebra) and if t is a matrix corepresentation then t is irreducible iff t' is irreducible. (For the proof, suppose that one of both corepresentations is not irreducible and bring it, by equivalence, in suitable block matrix form.)

Let $\mathcal{A}$ be a Hopf algebra. Because of (1.4), the mapping $\Delta: \mathcal{A} \rightarrow \mathcal{A} \otimes \mathcal{A}$ is a corepresentation of $\mathcal{A}$ on $\mathcal{A}$, the so-called *right regular corepresentation*. Similarly, the mapping $\tau \circ (S \otimes \text{id}) \circ \Delta: \mathcal{A} \rightarrow \mathcal{A} \otimes \mathcal{A}$ defines the *left regular corepresentation* on $\mathcal{A}$. This is indeed a corepresentation:

$$(\tau \circ (S \otimes \text{id}) \circ \Delta)(a) = \sum_{(a)} a_{(2)} \otimes S(a_{(1)}), \quad a \in \mathcal{A}.$$

Then both $(\tau \circ (S \otimes \text{id}) \circ \Delta) \otimes \text{id}$ and $\text{id} \otimes \Delta$ applied to the above right hand side yield

$$\sum_{(a)} a_{(3)} \otimes S(a_{(2)}) \otimes S(a_{(1)}).$$

Proposition 1.28 Let $\mathcal{A}$ be a Hopf algebra with invertible antipode. Let $\{t^\alpha\}_{\alpha \in I}$ (I some index set) be a collection of mutually inequivalent irreducible matrix corepresentations of $\mathcal{A}$. Then the set of all matrix elements t_{ij}^α is a set of linearly independent elements.

Proof Consider first a single irreducible d -dimensional matrix corepresentation t of $\mathcal{A}$. We have

$$\Delta(t_{ij}) = \sum_k t_{ik} \otimes t_{kj}, \quad (1.46)$$

$$(\tau \circ (S \otimes \text{id}) \circ \Delta)(t_{ij}) = \sum_k t_{kj} \otimes t'_{ki}. \quad (1.47)$$

Let $\tilde{\Delta}$ be the corepresentation of $\mathcal{A} \otimes \mathcal{A}$ on $\mathcal{A}$ which is defined by

$$\tilde{\Delta} := ((\tau \circ (S \otimes \text{id}) \circ \Delta) \otimes \text{id}) \circ \Delta: a \mapsto \sum_{(a)} S(a_{(2)}) \otimes a_{(1)} \otimes a_{(3)}: \mathcal{A} \rightarrow \mathcal{A} \otimes \mathcal{A} \otimes \mathcal{A}.$$

Then, by (1.46) and (1.47), we have

$$\tilde{\Delta}(t_{ij}) = \sum_{k,l} t_{kl} \otimes t'_{ki} \otimes t_{lj}.$$

Thus $\text{Span}\{t_{ij}\}$ is an invariant subspace of $\mathcal{A}$ with respect to $\tilde{\Delta}$. Denote the restriction of the corepresentation $\tilde{\Delta}$ to this subspace by $\hat{t}$.

Let $e_1, \dots, e_d$ be the standard basis of $\mathbb{C}^d$. We can consider both t and t' as corepresentations of $\mathcal{A}$ on $\mathbb{C}^d$, cf. (1.37). Let $\hat{t}$ be the corepresentation of $\mathcal{A} \otimes \mathcal{A}$ on $\mathbb{C}^d \otimes \mathbb{C}^d$ which is the tensor product of the corepresentations t' and t of $\mathcal{A}$ on $\mathbb{C}^d$. By Lemma 1.26 this corepresentation is irreducible. Then

$$\hat{t}(e_i \otimes e_j) = \sum_{k,l} (e_k \otimes e_l) \otimes t'_{ki} \otimes t_{lj}.$$

Thus the surjective linear mapping $A: \mathbb{C}^d \otimes \mathbb{C}^d \rightarrow \text{Span}\{t_{ij}\}$ defined by $A(e_i \otimes e_j) := t_{ij}$ is an intertwining operator for the corepresentations t and $\hat{t}$. Then $A^{-1}(0)$ is an invariant subspace of $\mathbb{C}^d \otimes \mathbb{C}^d$ with respect to $\hat{t}$, certainly a proper subspace, since otherwise $t_{11} = 0$, while $\varepsilon(t_{11}) = 1$. Because of the irreducibility of $\hat{t}$ we then have $A^{-1}(0) = \{0\}$. Thus A is bijective. Hence the t_{ij} are linearly independent.

Next consider m mutually inequivalent irreducible matrix corepresentations t^l ($l = 1, \dots, m$) of $\mathcal{A}$, where $t^l = (t_{ij}^l)_{i,j=1,\dots,d_l}$. Denote the linear span of all elements t_{ij}^l by V . Then, with the above notation, V is an invariant subspace of $\mathcal{A}$ with respect to the corepresentation $\tilde{\Delta}$ of $\mathcal{A} \otimes \mathcal{A}$ on $\mathcal{A}$. Let $W := \oplus_{l=1}^m W_l$, where $W_l := \mathbb{C}^{d_l} \otimes \mathbb{C}^{d_l}$. On W we have the corepresentation of $\mathcal{A} \otimes \mathcal{A}$ which is the direct

sum of the tensor product corepresentations of $(t^l)'$ with t^l and the W_l are invariant irreducible subspaces of W . The irreducible corepresentations of $\mathcal{A} \otimes \mathcal{A}$ on the W_l are mutually inequivalent.

Similarly as above we have a surjective intertwining operator $A: W \rightarrow V$ with respect to these two corepresentations of $\mathcal{A} \otimes \mathcal{A}$. Then $A^{-1}(0)$ is an invariant subspace of W . Suppose that $A^{-1}(0) \neq \{0\}$. Then we can take an invariant irreducible subspace U of $A^{-1}(0)$. By Lemma 1.25, $U = W_l$ for some l . This implies that, for this l , $t_{ij}^l = 0$, which contradicts that $\varepsilon(t_{11}^l) = 1$. $\square$

Lemma 1.29 Let $\mathcal{A}$ be a Hopf algebra with invertible antipode. Let s and t be irreducible matrix corepresentations of $\mathcal{A}$. Then s and t are equivalent iff $\text{Span}\{s_{ij}\} = \text{Span}\{t_{ij}\}$.

Proof Clearly, if s and t are equivalent then the two spans coincide. Conversely, let $V := \text{Span}\{s_{ij}\} = \text{Span}\{t_{ij}\}$. Then V is an invariant subspace of $\mathcal{A}$ with respect to the corepresentation Δ of $\mathcal{A}$ on $\mathcal{A}$. By Proposition 1.28 and formula (1.46) the restriction of Δ to V is a corepresentation of $\mathcal{A}$ which is a direct sum of d copies of s and also a direct sum of d copies of t . By Lemma 1.25 we conclude that s is equivalent to t . $\square$

Definition 1.30 Let s and t be corepresentations of a Hopf algebra $\mathcal{A}$ on vector spaces V and W , respectively. Then the *tensor product* $s \otimes t$ of s and t is defined as the corepresentation of $\mathcal{A}$ on $V \otimes W$ given by the mapping $(\text{id} \otimes \text{id} \otimes m) \circ (\text{id} \otimes \tau \otimes \text{id}) \circ (s \otimes t): V \otimes W \rightarrow (V \otimes W) \otimes \mathcal{A}$, i.e.,

$$(s \otimes t)(v \otimes w) = \sum_{(v), (w)} v_{(1)} \otimes w_{(1)} \otimes v_{(2)} w_{(2)}.$$

If s and t are matrix corepresentations then $s \otimes t$ can also be considered as a matrix corepresentation:

$$(s \otimes t)_{ij,kl} := s_{ik} t_{jl}.$$

Note that this tensor product is associative, but not commutative. The present tensor product should not be confused with the corepresentation of $\mathcal{A} \otimes \mathcal{A}$ obtained as a tensor product of s and t .

Definition 1.31 We call a corepresentation of a Hopf $*$ -algebra *unitarizable* if it is equivalent to a unitary corepresentation. In particular, a matrix corepresentation t of $\mathcal{A}$ is unitarizable iff there is an invertible complex square matrix A such that $s := AtA^{-1}$ satisfies $S(s_{ij}) = s_{ji}^*$.

Note that tensor products of unitary (resp. unitarizable) matrix corepresentations are again unitary (resp. unitarizable).

2. Compact quantum groups

This section gives a new approach to the theory of compact matrix quantum groups as developed by Woronowicz [9], [10], [11]. While Woronowicz's definition already involves C^* -algebras and his further development of the theory also heavily uses C^* -algebra theory, the present approach only uses Hopf $*$ -algebras, and mentions the connection with C^* -algebras at a much later stage, more as a side remark. This new approach was developed in cooperation with M. S. Dijkhuizen (CWI, Amsterdam). It will also be described in his forthcoming Ph.D. thesis [3]. The relationship with other work on compact quantum groups will be discussed at the end of this section. In particular, we acknowledge that Effros & Ruan [4] introduced CQG algebras (named differently by them) earlier. However, they developed their theory in a very different way.

2.1. CQG algebras and CMQG algebras. Let G be a compact group. Let $\Pi(G)$ be the linear span of all matrix elements of irreducible unitary (hence finite dimensional) representations of G . Then $\Pi(G)$ is a commutative Hopf $*$ -algebra with Hopf $*$ -algebra operations as described in Examples 1.4 and 1.11. We speak of a *compact matrix group* if, for some n , G is (isomorphic to) a closed subgroup of the group $U(n)$ of $n \times n$ unitary matrices. For a compact group G there are the following equivalent statements: G is a compact matrix group iff G is a compact Lie group iff $\Pi(G)$ is finitely generated. This suggests the following definitions.

Definition 2.1 A Hopf $*$ -algebra $\mathcal{A}$ is said to be associated with a *compact quantum group* if it is the linear span of the matrix elements of the unitary (finite dimensional) matrix corepresentations of $\mathcal{A}$. Then $\mathcal{A}$ is called a *CQG algebra*.

Because of Proposition 1.23(b), a CQG algebra $\mathcal{A}$ is also the linear span of the matrix elements of its irreducible unitary matrix corepresentations. Denote by $\hat{\mathcal{A}}$ the collection of all equivalence classes of irreducible unitary matrix corepresentations of $\mathcal{A}$. Choose, for each $\alpha \in \hat{\mathcal{A}}$, a unitary matrix corepresentation $(t_{ij}^\alpha)_{i,j=1,\dots,d_\alpha}$ belonging to class α . Then, by Proposition 1.28, the set of all t_{ij}^α forms a basis of $\mathcal{A}$. Write

$$\mathcal{A}_\alpha := \text{Span}\{t_{ij}^\alpha\}_{i,j=1,\dots,d_\alpha}, \quad \alpha \in \hat{\mathcal{A}}.$$

Then $\mathcal{A}$ is the direct sum of the $\mathcal{A}_\alpha$. Let 1 be the element of $\hat{\mathcal{A}}$ for which t^1 is the one-dimensional matrix corepresentation (1). For $\alpha \in \hat{\mathcal{A}}$ let α' be the element of $\hat{\mathcal{A}}$ such that $(t^\alpha)'$ is equivalent to $t^{\alpha'}$.

Proposition 2.2 Let $\mathcal{A}$ be a CQG algebra and let the t^α be as above.

- (a) Each irreducible matrix corepresentation of $\mathcal{A}$ is equivalent to some t^α ($\alpha \in \hat{\mathcal{A}}$).
- (b) Each irreducible matrix corepresentation of $\mathcal{A}$ is unitarizable.
- (c) If s is a unitarizable matrix corepresentation of $\mathcal{A}$ then so is s' .

Proof (b) immediately follows from (a). For (c) observe that a unitarizable s is a direct sum of irreducible corepresentations. Hence s' is also a direct sum of irreducible corepresentations. Then apply (b). Thus we have to prove (a). Let s be an irreducible matrix corepresentation of $\mathcal{A}$. The spaces $\mathcal{A}_\alpha$ are invariant irreducible subspaces of $\mathcal{A}$ with respect to the corepresentation of $\mathcal{A} \otimes \mathcal{A}$ on $\mathcal{A}$ which is the tensor product of the left regular and the right regular corepresentation (cf. the proof of Proposition 1.28), and so is $V := \text{Span}\{s_{ij}\}$. Moreover, the corepresentations obtained by restriction to the various $\mathcal{A}_\alpha$ are mutually inequivalent. By Lemma 1.25 $V = \mathcal{A}_\alpha$ for some $\alpha \in \widehat{\mathcal{A}}$. By Lemma 1.29 we see that s is equivalent to t^α . $\square$

Definition 2.3 A Hopf $*$ -algebra $\mathcal{A}$ is said to be associated with a *compact matrix quantum group* if it is a finitely generated CQG algebra. Then $\mathcal{A}$ is called a *CMQG algebra*.

Proposition 2.4 For a Hopf $*$ -algebra the following conditions are equivalent.

- (i) $\mathcal{A}$ is a CMQG algebra.
- (ii) There is a unitary matrix corepresentation v of $\mathcal{A}$ such that $\mathcal{A}$ is generated as an algebra by the matrix coefficients of v .
- (iii) There is a matrix corepresentation u of $\mathcal{A}$ such that both u and u' are unitarizable and $\mathcal{A}$ as an algebra is generated by the matrix elements of u and u' .

Proof

(i) $\Rightarrow$ (ii) $\Rightarrow$ (iii): Assume (i). Each of the generators is a linear combination of matrix elements of unitary matrix corepresentations of $\mathcal{A}$. Hence, by taking direct sums, there is a unitary matrix corepresentation u of $\mathcal{A}$ of which the matrix elements generate $\mathcal{A}$. Then u' is unitarizable by Proposition 2.2.

(iii) $\Rightarrow$ (ii) $\Rightarrow$ (i): Assume (iii). Let v be the direct sum of two unitary matrix corepresentations which are equivalent to u respectively u' . Then $\mathcal{A}$ is generated by the matrix elements of v . Each product of matrix elements of v is a matrix element of some multiple tensor product of v . Such tensor products are again unitary corepresentations. Thus $\mathcal{A}$ is a finitely generated CQG algebra. $\square$

2.2. The Haar functional. Define on a CQG algebra $\mathcal{A}$ the *Haar functional* h as the linear mapping $h: \mathcal{A} \rightarrow \mathbb{C}$ such that

$$h(a) := \begin{cases} 0 & \text{if } a \in \mathcal{A}_\alpha, \alpha \neq 1, \\ 1 & \text{if } a = 1, \end{cases} \quad (2.1)$$

The functional h satisfies

$$(h \otimes \text{id})(\Delta(a)) = h(a) 1 = (\text{id} \otimes h)(\Delta(a)), \quad a \in \mathcal{A}. \quad (2.2)$$

Indeed, if a equals some basis element t_{ij}^α of $\mathcal{A}$ then (2.2) takes the form

$$\sum_k h(t_{ik}^\alpha) t_{kj}^\alpha = h(t_{ij}^\alpha) 1 = \sum_k h(t_{kj}^\alpha) t_{ik}^\alpha, \quad (2.3)$$

and these identities are indeed implied by (2.1).

On the other hand, if h would be any linear functional on $\mathcal{A}$ satisfying the first (or the second) identity in (2.2) then we obtain from (2.3) by linear independence of the t_{ij}^α that $h(t_{ij}^\alpha) = 0$ for $\alpha \neq 1$. Thus, h is determined up to a constant factor by each of the identities in (2.2) and h can next be normalized by

$$h(1) = 1. \quad (2.4)$$

Since $S(\mathcal{A}_\alpha) = \mathcal{A}_{\alpha'}$, (2.1) implies that

$$h(S(a)) = h(a), \quad a \in \mathcal{A}. \quad (2.5)$$

If G is a compact group then the normalized Haar measure, as a linear functional on $\Pi(G)$, can be interpreted as a Haar functional. Indeed,

$$h(a) := \int_G a(x) dx, \quad a \in \Pi(G),$$

and

$$((h \otimes \text{id})(\Delta(a)))(y) = \int_G a(xy) dx = \int_G a(x) dx = h(a) 1(y),$$

and similarly for the other identity. In this case we also have that $h(aa^*) = \int_G |a(x)|^2 dx > 0$ for $a \neq 0$. We want to prove this positivity result also for h in the case of a CQG algebra.

The following Lemma will be crucial. For the moment we will assume that $\mathcal{A}$ is a Hopf algebra with invertible antipode (not necessarily a Hopf $*$ -algebra) and that h is a linear functional on $\mathcal{A}$ satisfying (2.2), (2.4) and (2.5).

Lemma 2.5 Let r and s be matrix corepresentations of $\mathcal{A}$. Then

$$\sum_l h(s_{ij} S(r_{kl})) r_{lm} = \sum_l s_{il} h(s_{lj} S(r_{km})), \quad (2.6)$$

$$\sum_l h(S(r_{ij}) s_{kl}) s_{lm} = \sum_l r_{il} h(S(r_{lj}) s_{km}). \quad (2.7)$$

With the notation

$$A_{il}^{(j,k)} := h(s_{ij} S(r_{kl})), \quad B_{il}^{(j,k)} := h(S(r_{ij}) s_{kl}), \quad (2.8)$$

the identities (2.6) and (2.7) can be rewritten as

$$A^{(j,k)} r = s A^{(j,k)}, \quad B^{(j,k)} s = r B^{(j,k)}. \quad (2.9)$$

Thus $A^{(j,k)}$ is an intertwining operator for r and s and $B^{(j,k)}$ is an intertwining operator for s and r .

Proof For the proof of (2.6) write

$$\begin{aligned} h(s_{ij} S(r_{kl})) 1 &= (\text{id} \otimes h)(\Delta(s_{ij} S(r_{kl}))) \\ &= \sum_{p,n} (\text{id} \otimes h)((s_{ip} \otimes s_{pj})(S(r_{nl}) \otimes S(r_{kn}))) = \sum_{p,n} h(s_{pj} S(r_{kn})) s_{ip} S(r_{nl}). \end{aligned}$$

Substitute this equality in the left hand side of (2.6) and next use that $\sum_l S(r_{nl}) r_{lm} = \delta_{nm} 1$. This settles (2.6). Similarly, (2.7) is obtained from a substitution of

$$h(S(r_{lj}) s_{km}) 1 = (h \otimes \text{id})(\Delta(S(r_{lj}) s_{km}))$$

in the right hand side of (2.7). $\square$

Let us assume that r and s are irreducible matrix corepresentations of $\mathcal{A}$. We will consider three special cases of Lemma 2.5: (i) r and s are inequivalent, (ii) $s=r$, (iii) $s = r''$ (i.e., $s_{ij} = S^2(r_{ij})$). For each case we apply one of the quantum Schur lemmas (Lemma 1.24(b),(c)).

Case (i) If r and s are inequivalent then (2.9) yields that $A_{il}^{(j,k)} = 0$ and $B_{il}^{(j,k)} = 0$. Hence, by (2.8),

$$h(s_{ij} S(r_{kl})) = 0, \quad h(S(r_{ij}) s_{kl}) = 0.$$

Case (ii) If $s = r$ then (2.9) yields that there are complex constants α_{jk} and β_{jk} such that $A^{(j,k)} = \alpha_{jk} I$ and $B_{il}^{(j,k)} = \beta_{jk} I$. Hence, by (2.8)

$$A_{il}^{(j,k)} = h(r_{ij} S(r_{kl})) = \alpha_{jk} \delta_{il}, \quad (2.10)$$

$$B_{il}^{(j,k)} = h(S(r_{ij}) r_{kl}) = \beta_{jk} \delta_{il}. \quad (2.11)$$

Moreover, if we fix $i = l$ and sum over $j = k$ then we obtain

$$\sum_j \alpha_{jj} = 1 = \sum_j \beta_{jj}. \quad (2.12)$$

Case (iii) If $s = r''$ then (2.6) and (2.7) together with (2.5) yield

$$\begin{aligned} \sum_l h(r_{kl} S(r_{ij})) r_{lm} &= \sum_l S^2(r_{il}) h(r_{km} S(r_{lj})), \\ \sum_l h(S(r_{kl}) r_{ij}) S^2(r_{lm}) &= \sum_l r_{il} h(S(r_{km}) r_{lj}). \end{aligned}$$

Put

$$\tilde{A}_{il}^{(j,k)} := h(r_{kl} S(r_{ij})) = A_{kj}^{(l,i)} = \alpha_{li} \delta_{kj}, \quad (2.13)$$

$$\tilde{B}_{il}^{(j,k)} := h(S(r_{kl}) r_{ij}) = B_{kj}^{(l,i)} = \beta_{li} \delta_{kj}, \quad (2.14)$$

where we refer to (2.10), (2.11). Then $\tilde{A}^{(j,k)} r = r'' \tilde{A}^{(j,k)}$ and $\tilde{B}^{(j,k)} r'' = r \tilde{B}^{(j,k)}$. Moreover, when we sum the first equality in (2.13) and in (2.14) over $i = l$ then we obtain that

$$\text{tr } \tilde{A}^{(j,k)} = \delta_{jk}, \quad \text{tr } \tilde{B}^{(j,k)} = \delta_{jk}.$$

Hence, there exists a nonzero intertwining operator F for r and r'' . Thus r and r'' are equivalent irreducible corepresentations (since S is assumed to be invertible, the irreducibility of r implies the irreducibility of r''), and F is an invertible operator, unique up to a constant factor and satisfying $\text{tr } F \neq 0$ and $\text{tr } F^{-1} \neq 0$. Hence there are complex constants $\tilde{\alpha}_{jk}$ and $\tilde{\beta}_{jk}$ such that

$$\begin{aligned} \tilde{A}_{il}^{(j,k)} &= \tilde{\alpha}_{jk} F_{il} = \alpha_{li} \delta_{kj}, \\ \tilde{B}_{il}^{(j,k)} &= \tilde{\beta}_{jk} (F^{-1})_{il} = \beta_{li} \delta_{kj}, \end{aligned}$$

where, in each line, the second equality follows from (2.13). Combination with (2.12) yields that

$$\tilde{\alpha}_{jk} \text{tr } F = \delta_{kj}, \quad \tilde{\beta}_{jk} \text{tr } F^{-1} = \delta_{kj}.$$

Hence

$$\tilde{A}_{il}^{(j,k)} = \delta_{kj} \frac{F_{il}}{\text{tr } F}, \quad \tilde{B}_{il}^{(j,k)} = \delta_{kj} \frac{(F^{-1})_{il}}{\text{tr } F^{-1}}. \quad (2.15)$$

Let us summarize the obtained results in the following Proposition.

Proposition 2.6 Let $\mathcal{A}$ be a Hopf algebra with invertible antipode and let h be a linear functional on $\mathcal{A}$ satisfying (2.2), (2.4) and (2.5). Let r be an irreducible matrix corepresentation of $\mathcal{A}$.

- (a) If s is an irreducible matrix corepresentation of $\mathcal{A}$ which is not equivalent to r then

$$h(s_{kl} S(r_{ij})) = 0, \quad h(S(r_{kl}) s_{ij}) = 0. \quad (2.16)$$

- (b) r is equivalent to r'' . Let F be an invertible intertwining operator for r and r'' . Then $\text{tr } F \neq 0$ and $\text{tr } F^{-1} \neq 0$ and

$$h(r_{kl} S(r_{ij})) = \delta_{kj} \frac{F_{il}}{\text{tr } F}, \quad (2.17)$$

$$h(S(r_{kl}) r_{ij}) = \delta_{kj} \frac{(F^{-1})_{il}}{\text{tr } F^{-1}}. \quad (2.18)$$

Proof (2.17) and (2.18) were obtained from (2.13), (2.14) and (2.15). $\square$

Now we assume that $\mathcal{A}$ is a CQG algebra.

Proposition 2.7 Let r be an irreducible unitary matrix corepresentation of the CQG algebra $\mathcal{A}$. Let the matrix F be an invertible intertwining operator for r and r'' as in Proposition 2.6. Then F is a constant multiple of a positive definite matrix. It can be uniquely normalized such that $\text{tr } F = \text{tr } F^{-1} > 0$.

Proof By Proposition 2.2(c) there is a unitary matrix corepresentation s such that r' is equivalent to s . Thus there is an invertible complex matrix A such that $sA = Ar'$. Let $\overline{A}$ be the matrix for which $(\overline{A})_{ij} := \overline{A_{ij}}$ and let tA be the transpose of A . Since $s'_{ij} = (s_{ij})^* = S(s_{ji})$ and $r'_{ij} = (r_{ij})^*$, $r''_{ij} = S(r'_{ji})$, we conclude from $sA = Ar'$ that $s'\overline{A} = \overline{A}r$ and ${}^tA s' = r'' {}^tA$. Hence ${}^tA \overline{A} r = r'' {}^tA \overline{A}$. Thus $F = \text{const. } {}^tA \overline{A}$ and ${}^tA \overline{A}$ is positive definite. $\square$

Theorem 2.8 The Haar functional on a CQG algebra satisfies

$$h(aa^*) > 0 \quad \text{if } a \neq 0,$$

Proof Propositions 2.6 and 2.7 imply that

$$h(t_{kl}^\alpha (t_{ij}^\beta)^*) = \delta_{\alpha\beta} \delta_{ki} G_{lj}^\alpha, \quad \alpha, \beta \in \widehat{\mathcal{A}},$$

for certain positive definite matrices G^α . Let $a := \sum_{\alpha, k, l} c_{kl}^\alpha t_{kl}^\alpha$ be an arbitrary element of $\mathcal{A}$. Then

$$h(aa^*) = \sum_{\alpha, k} \sum_{l, j} c_{kl}^\alpha \overline{c_{kj}^\alpha} G_{lj}^\alpha \geq 0.$$

If $h(aa^*) = 0$ then $\sum_{l, j} c_{kl}^\alpha \overline{c_{kj}^\alpha} G_{lj}^\alpha = 0$ for all α and k . By positive definiteness of the G^α this implies that all coefficients c_{kl}^α are 0, i.e., $a = 0$. $\square$

Corollary 2.9 Every finite dimensional corepresentation of a CQG algebra $\mathcal{A}$ is unitarizable, and hence decomposable as a direct sum of irreducible finite dimensional corepresentations.

Proof Let r be a corepresentation of $\mathcal{A}$ on a vector space V with some inner product $(\cdot, \cdot)$. Define a new sesquilinear form on V by

$$\langle v, w \rangle := \sum_{(v), (w)} (v_{(1)}, w_{(1)}) h(w_{(2)}^* v_{(2)}).$$

Then r is unitary in the sense of (1.41) with respect to this sesquilinear form. Indeed,

$$\begin{aligned} \sum_{(v), (w)} \langle v_{(1)}, w_{(1)} \rangle w_{(2)}^* v_{(2)} &= \sum_{(v), (w)} (v_{(1)}, w_{(1)}) h(w_{(2)}^* v_{(2)}) w_{(3)}^* v_{(3)} \\ &= \sum_{(v), (w)} (v_{(1)}, w_{(1)}) h(w_{(2)}^* v_{(2)}) 1 = \langle v, w \rangle 1. \end{aligned}$$

It is left to show that $\langle \cdot, \cdot \rangle$ is positive definite. Let $e_1, \dots, e_n$ be an orthonormal basis of V with respect to the inner product $(\cdot, \cdot)$ and write r as a matrix corepresentation with respect to this basis. Then

$$\langle e_i, e_j \rangle = \sum_{k, l} (e_k, e_l) h(r_{lj}^* r_{ki}) = \sum_k h(r_{kj}^* r_{ki}).$$

Hence

$$\langle \sum_i c_i e_i, \sum_j c_j e_j \rangle = \sum_k h((\sum_j c_j r_{kj})^* (\sum_i c_i r_{ki})) \geq 0,$$

and if the left hand side equals 0 then $\sum_i c_i r_{ki} = 0$ for all k , hence $c_k = \varepsilon(\sum_i c_i r_{ki}) = 0$ for all k . $\square$

2.3. The C^* -completion of a CQG algebra. Recall that a C^* -algebra is a Banach algebra A which is also a $*$ -algebra such that $\|aa^*\| = \|a\|^2$ for all $a \in A$. In particular, the space $\mathcal{L}(\mathcal{H})$ of all bounded linear operators on a Hilbert space $\mathcal{H}$ forms a C^* -algebra with identity. By a $*$ -representation of a $*$ -algebra $\mathcal{A}$ on a Hilbert space $\mathcal{H}$ we mean a $*$ -algebra homomorphism $\pi: \mathcal{A} \rightarrow \mathcal{L}(\mathcal{H})$.

On a compact group G the $*$ -algebra $\Pi(G)$ can be equipped with the sup norm. The space $C(G)$ of all continuous functions on G is the completion of $\Pi(G)$ with respect to this norm and $C(G)$ is a commutative C^* -algebra with identity. Both for the $*$ -algebra $\Pi(G)$ and for the C^* -algebra $C(G)$ the irreducible $*$ -representations on a Hilbert space are precisely the one-dimensional $*$ -homomorphisms $a \mapsto a(x)$ ($x \in G$).

This suggests the following strategy for the construction of an analogue of $C(G)$ for a CQG algebra $\mathcal{A}$:

- 1) Define a seminorm

$$\|a\| := \sup_{\pi} \|\pi(a)\| \quad (2.19)$$

on $\mathcal{A}$, where the supremum runs over all $*$ -representations π of the $*$ -algebra $\mathcal{A}$.

- 2) Show that this seminorm is a norm.
- 3) Let A be the completion of $\mathcal{A}$ with respect to this norm.

First observe that $\|a\|$ defined by (2.19) is finite for all $a \in \mathcal{A}$. Indeed, if a is expanded in terms of the basis of $\mathcal{A}$ by the finite sum

$$a = \sum_{\alpha, i, j} c_{ij}^{\alpha} t_{ij}^{\alpha} \quad (2.20)$$

then $\|a\| \leq \sum_{\alpha, i, j} |c_{ij}^{\alpha}| \|t_{ij}^{\alpha}\|$. Now use the following lemma.

Lemma 2.10 Let r be a unitary matrix corepresentation of a CQG algebra $\mathcal{A}$. Let V be a complex vector space with hermitian inner product and let $|v| := (v, v)^{\frac{1}{2}}$ define the norm on V . Let π be an algebra homomorphism from $\mathcal{A}$ to the algebra of all linear operators on V such that $(\pi(a)v, w) = (v, \pi(a^*)w)$ for all $a \in \mathcal{A}$ and all $v, w \in V$. Then $\pi(r_{ij})$ is a bounded linear operator on V of norm ≤ 1 for all i, j . In particular, π uniquely extends to a $*$ -algebra representation of $\mathcal{A}$ on the Hilbert space completion of V .

Proof Since r is unitary, we have $\sum_k r_{kj}^* r_{kj} = 1$. Hence, for all $v \in V$,

$$\|v\|^2 = (v, v) = \sum_k (\pi(r_{kj}^* r_{kj}) v, v) = \sum_k (\pi(r_{kj}) v, \pi(r_{kj}) v) \geq \|\pi(r_{ij}) v\|^2. \quad \square$$

One consequence of this Lemma is that, for $a \in \mathcal{A}$ with expansion (2.20), all norms $\|\pi(a)\|$, and hence $\|a\|$, are bounded by $\sum_{\alpha, i, j} |c_{ij}^{\alpha}|$.

Another consequence concerns the left regular representation of $\mathcal{A}$ on $\mathcal{A}$, which is defined by $\lambda(a)b := ab$. This is an algebra representation, which is *faithful*, i.e.,

if $\lambda(a) = 0$ then $a = 0$. Now equip $\mathcal{A}$ with the inner product $(a, b) := h(b^*a)$. (By the properties of h this is indeed an inner product.) Clearly, $(\lambda(a)b, c) = (b, \lambda(a^*)c)$. Since each $a \in \mathcal{A}$ is a linear combination of matrix elements of unitary corepresentations, it follows by Lemma 2.10 that each $\lambda(a)$ is a bounded linear operator on the inner product space $\mathcal{A}$. Hence λ extends to a $*$ -representation of $\mathcal{A}$ on the Hilbert space completion of this inner product space. (I thank P. Podleś for this observation.) By faithfulness it follows that the seminorm (2.19) is a norm.

Because $\|\pi(ab)\| \leq \|\pi(a)\| \|\pi(b)\|$ and $\|\pi(a)\|^2 = \|\pi(aa^*)\|$ for each $*$ -representation π of $\mathcal{A}$, it follows that the norm (2.19) is a C^* -norm, i.e., $\|ab\| \leq \|a\| \|b\|$ and $\|aa^*\| = \|a\|^2$. Thus the norm completion A of $\mathcal{A}$ is a well-defined C^* -algebra with identity.

Since, for each element a of a C^* -algebra A , there exists an irreducible $*$ -representation π of A for which $\|\pi(a)\| = \|a\|$ (cf. for instance Arveson [2, Corollary to Theorem 1.7.2]), the norm definition (2.19) does not change when we take the supremum there only over the irreducible $*$ -representations of $\mathcal{A}$.

The counit and comultiplication on $\mathcal{A}$ have unique continuous extensions to A . For the counit this is obvious, since ε is a one-dimensional $*$ -representation of $\mathcal{A}$, so $|\varepsilon(a)| \leq \|a\|$.

For the C^* -extension of Δ we need a suitable C^* -norm on the algebraic tensor product $A \otimes A$. We choose the *injective cross norm*, i.e., for $a \in A \otimes A$, the norm $\|a\|$ is defined as the supremum of the numbers $\|(\pi_1 \otimes \pi_2)(a)\|$, where π_1 and π_2 run over the $*$ -representations of the C^* -algebra A . Then Δ continuously extends to a mapping from A to the completion of $A \otimes A$ with respect to this norm, since $a \mapsto (\pi_1 \otimes \pi_2)(\Delta(a))$ is a $*$ -representation of $\mathcal{A}$ for any two $*$ -representations π_1 and π_2 of $\mathcal{A}$, so $\|(\pi_1 \otimes \pi_2)(\Delta(a))\| \leq \|a\|$. (The $*$ -representation $a \mapsto (\pi_1 \otimes \pi_2)(\Delta(a))$ of $\mathcal{A}$ is called the *tensor product* $\pi_1 \otimes \pi_2$ of the $*$ -representations π_1 and π_2 , cf. Remark 1.8.)

2.4. A class of multiplicative linear functionals. Let $\mathcal{A}$ be a CQG algebra. According to Proposition 2.7 we can choose for each $\alpha \in \hat{\mathcal{A}}$ a positive definite complex matrix F_α which is an intertwining operator for the matrix corepresentations t^α and $(t^\alpha)''$ and which is uniquely normalized such that $\text{tr } F_\alpha = \text{tr } (F_\alpha)^{-1}$. We can now rewrite (2.16), (2.17) and (2.18) as

$$h(t_{kl}^\beta (t_{ji}^\alpha)^*) = \delta_{\alpha\beta} \delta_{kj} \frac{(F_\alpha)_{il}}{\text{tr } F_\alpha}, \quad (2.21)$$

$$h((t_{lk}^\beta)^* t_{ij}^\alpha) = \delta_{\alpha\beta} \delta_{kj} \frac{(F_\alpha^{-1})_{il}}{\text{tr } F_\alpha^{-1}}. \quad (2.22)$$

These can be considered as the analogues for CQG algebras of Schur's orthogonality relations. Woronowicz, for the case of his compact matrix quantum groups, gave these quantum Schur orthogonality relations in [9, (5.14), (5.15)].

The remainder of this subsection is an account of Theorem 5.6 in Woronowicz [9]. It describes the so-called modular properties of the Haar functional h . Write F_α

as

$$F_\alpha = U \begin{pmatrix} \lambda_1 & & 0 \\ & \ddots & \\ 0 & & \lambda_{d_\alpha} \end{pmatrix} U^{-1},$$

where U is a unitary matrix and $\lambda_1, \dots, \lambda_{d_\alpha} > 0$. Now define complex powers F_α^z ($z \in \mathbb{C}$) of F_α by

$$F_\alpha^z := U \begin{pmatrix} \lambda_1^z & & 0 \\ & \ddots & \\ 0 & & \lambda_{d_\alpha}^z \end{pmatrix} U^{-1}.$$

Then the matrix elements $(F_\alpha^z)_{ij}$ are entire analytic functions of z and there are constants $M > 0$ and $\mu \in \mathbb{R}$ such that

$$|(F_\alpha^z)_{ij}| \leq M e^{\mu \operatorname{Re} z} \quad \text{for all } z \in \mathbb{C}.$$

Note also that $F_\alpha^z F_\alpha^{z'} = F_\alpha^{z+z'}$.

Remark 2.11 If f is an entire analytic function satisfying the estimate $|f(z)| \leq M e^{\mu \operatorname{Re} z}$ for some $M > 0$ and $\mu \in \mathbb{R}$ then we can find a real constant c such that the function $g(z) := e^{cz} f(z)$ satisfies the estimate

$$\exists M > 0 \quad \exists \mu \in (0, \pi) \quad \operatorname{Re} z \geq 0 \implies |g(z)| \leq M e^{\mu |z|}. \quad (2.23)$$

Carlson's theorem (cf. for instance Titchmarsh [7, §5.81] states that, if a holomorphic function g on $\{z \mid \operatorname{Re} z \geq 0\}$ satisfies (2.23) and if moreover $g(1) = g(2) = \dots = 0$, then g will vanish identically. Hence, if the above function f will vanish on the set $\{1, 2, \dots\}$ then it will also vanish identically.

Now define, for each $z \in \mathbb{C}$, the element $f_z \in \mathcal{A}^*$ by specifying it on our linear basis of $\mathcal{A}$:

$$f_z(t_{ij}^\alpha) := (F_\alpha^z)_{ij}.$$

Note that this definition is independent of the choice of the matrix corepresentation t^α in the class $\alpha \in \overline{\mathcal{A}}$. The following properties of the f_z quickly follow from the definition of f_z .

Proposition 2.12

- (a) $f_1(t_{ij}^\alpha) = (F_\alpha)_{ij}$, $f_{-1}(t_{ij}^\alpha) = ((F_\alpha)^{-1})_{ij}$.
- (b) $f_0 = \varepsilon$.
- (c) $f_z(1) = 1$.
- (d) $f_z f_{z'} = f_{z+z'}$.
- (e) For each $a \in \mathcal{A}$ the function $z \mapsto f_z(a)$ is an entire analytic function and there are constants $M > 0$ and $\mu \in \mathbb{R}$ such that $|f_z(a)| \leq M e^{\mu \operatorname{Re} z}$.

For some further properties we will give the proof.

Proposition 2.13 $S^2(a) = f_{-1}.a.f_1$.

Proof It follows from Proposition 2.7 that

$$\begin{aligned}
 S^2(t_{ij}^\alpha) &= \sum_{k,l} (F_\alpha)_{ik} t_{kl}^\alpha (F_\alpha^{-1})_{lj} \\
 &= (f_1 \otimes \text{id} \otimes f_{-1})((\Delta \otimes \text{id}) \Delta(t_{ij}^\alpha)) \\
 &= f_{-1} \cdot t_{ij}^\alpha \cdot f_1. \quad \square
 \end{aligned}$$

Proposition 2.14 If $a, b \in \mathcal{A}$ then

$$h(ab) = h(b(f_1 \cdot a \cdot f_1)). \quad (2.24)$$

Proof Let $\alpha, \beta \in \widehat{\mathcal{A}}$. It is enough to verify (2.24) for $a := t_{kl}^\beta$ and $b := (t_{ij}^\alpha)^*$. We apply (2.21) and (2.22). If $\alpha \neq \beta$ then both sides of (2.24) are 0. If $\alpha = \beta$ then

$$\begin{aligned}
 h((t_{ij}^\alpha)^* (f_1 \cdot t_{kl}^\alpha \cdot f_1)) &= \sum_{m,n} (F_\alpha)_{km} (F_\alpha)_{nl} h((t_{ij}^\alpha)^* t_{mn}^\alpha) \\
 &= \sum_{m,n} (F_\alpha)_{km} (F_\alpha)_{nl} \frac{\delta_{jn} (F_\alpha^{-1})_{mi}}{\text{tr } F_\alpha^{-1}} \\
 &= \frac{\delta_{ik} (F_\alpha)_{jl}}{\text{tr } F_\alpha^{-1}} = \frac{\delta_{ik} (F_\alpha)_{jl}}{\text{tr } F_\alpha} = h(t_{kl}^\alpha (t_{ij}^\alpha)^*),
 \end{aligned}$$

where we also used the chosen normalization for F_α . $\square$

The last result justifies the normalization of the F_α . Note on the other hand that Propositions 2.13 and 2.14, together with the property $f_1 f_{-1} = \varepsilon = f_{-1} f_1$ completely determine f_1 and f_{-1} as elements of $\mathcal{A}^*$. The other f_z are then determined by Proposition 2.12(d),(e), in view of Remark 2.11.

A linear functional f on an algebra $\mathcal{A}$ is called *central* if $f(ab) = f(ba)$ for all $a, b \in \mathcal{A}$. Thus h is not necessarily central, but its non-centrality is well controlled by (2.24).

Lemma 2.15 For $a, b \in \mathcal{A}$ we have

$$f_1 \cdot (ab) \cdot f_1 = (f_1 \cdot a \cdot f_1) (f_1 \cdot b \cdot f_1). \quad (2.25)$$

Proof Apply (2.24):

$$h(c(f_1 \cdot (ab) \cdot f_1)) = h(abc) = h(bc(f_1 \cdot a \cdot f_1)) = h(c(f_1 \cdot a \cdot f_1)(f_1 \cdot b \cdot f_1)).$$

Now use that $d \in \mathcal{A}$ equals 0 if $h(cd) = 0$ for all $c \in \mathcal{A}$. $\square$

A linear functional $f \in \mathcal{A}^*$ is called *multiplicative* if $f(ab) = f(a)f(b)$ and $f(1) = 1$. The multiplicative linear functionals form again an algebra. (This is true for $\mathcal{A}$ an arbitrary bialgebra.) Indeed, if $f, g \in \mathcal{A}$ are multiplicative then

$$\begin{aligned} (fg)(ab) &= \sum_{(a),(b)} f(a_{(1)}b_{(1)}) g(a_{(2)}b_{(2)}) \\ &= \sum_{(a),(b)} f(a_{(1)}) g(a_{(2)}) f(b_{(1)}) g(b_{(2)}) = (fg)(a) (fg)(b). \end{aligned}$$

Proposition 2.16 For all $z \in \mathbb{C}$ f_z is a multiplicative linear functional:

$$f_z(ab) = f_z(a) f_z(b), \quad a, b \in \mathcal{A}. \quad (2.26)$$

Proof Apply ε to both sides of (2.25). Then we get $f_1^2(ab) = f_1^2(a) f_1^2(b)$. Hence (2.26) is true for $z = 2$. By the above remark, (2.26) is then true for $z = 2, 4, 6, \dots$. Now apply Proposition 2.12(e) and Remark 2.11. $\square$

Proposition 2.16 tells us in particular that f_1 is multiplicative. I wonder if a more algebraic proof, not using Carlson's theorem, is possible for this fact.

Remark 2.17 As a consequence of this Proposition, f_1 and f_{-1} are already determined in a CMQG algebra by their values on the matrix elements v_{ij} , where v is as in Proposition 2.4(ii). This also determines the right hand sides of (2.21) and (2.22), since $(F_\alpha^{\pm 1})_{ij} = f_{\pm 1}(t_{ij}^\alpha)$.

Let us finally consider expressions for $S(f_z)$ and $(f_z)^*$. Note that a nonzero multiplicative linear functional $f \in \mathcal{A}^*$ has two-sided inverse $S(f)$. (This is true for $\mathcal{A}$ an arbitrary Hopf algebra.) Indeed,

$$\begin{aligned} \varepsilon(a) &= f(\varepsilon(a)1) = f(m \circ (S \otimes \text{id})\Delta(a)) \\ &= (f \otimes f)((S \otimes \text{id})\Delta(a)) = (S(f) \otimes f)(\Delta(a)) = (S(f)f)(a). \end{aligned}$$

This shows that

$$S(f_z) = f_{-z}. \quad (2.27)$$

In order to compute $(f_z)^*$ observe that

$$(f_z)^*(t_{ij}^\alpha) = \overline{f_z((S(t_{ij}^\alpha))^*)} = \overline{f_z(t_{ji}^\alpha)} = \overline{(F_\alpha^z)_{ji}} = (F_\alpha^{\bar{z}})_{ij} = f_{\bar{z}}(t_{ij}^\alpha).$$

Hence

$$(f_z)^* = f_{\bar{z}}. \quad (2.28)$$

2.5. Comparison with other literature

(a) Woronowicz [9], [10], [11]

Woronowicz, in his influential 1987 paper [9], gives the following definition of a *compact matrix quantum group* (originally called *compact matrix pseudogroup*). It is a pair (A, u) , where A is a unital C^* -algebra and $u = (u_{ij})_{i,j=1,\dots,N}$ is an $N \times N$ matrix with entries in A , such that the following properties hold.

- 1) The unital $*$ -subalgebra $\mathcal{A}$ of A generated by the entries of u is dense in A .
- 2) There exists a (necessarily unique) C^* -homomorphism $\Delta: A \rightarrow A \otimes A$ such that $\Delta(u_{ij}) = \sum_{k=1}^N u_{ik} \otimes u_{kj}$.
- 3) There exists a (necessarily unique) linear antimultiplicative mapping $S: \mathcal{A} \rightarrow \mathcal{A}$ such that $S \circ * \circ S \circ * = \text{id}$ on $\mathcal{A}$ and $\sum_{k=1}^N S(u_{ik}) u_{kj} = \delta_{ij} 1 = \sum_{k=1}^N u_{ik} S(u_{kj})$.

In his note [11] Woronowicz shows that, instead of 3), we may equivalently require:

- 3') The matrix u and its transpose are invertible.

Woronowicz now essentially shows (cf. [9, Prop. 1.8]) that there exists a (necessarily unique) $*$ -homomorphism $\varepsilon: \mathcal{A} \rightarrow \mathbb{C}$ such that $\varepsilon(u_{ij}) = \delta_{ij}$ and that $\mathcal{A}$ becomes a Hopf $*$ -algebra with comultiplication Δ , counit ε and antipode S . In [9] the notation Φ, e, κ is used instead of our Δ, ε, S , respectively. Note that the above $*$ -algebra $\mathcal{A}$ is very close to what we have defined as a CMQG algebra (cf. Proposition 2.2). However, it is not postulated and not yet obvious in the beginning of [9] that the corepresentations u and u' are unitarizable.

A central result in the paper (see [9, Theorem 4.2]) is the existence of a *state* (normalized positive linear functional) h on the C^* -algebra A such that $(h \otimes \text{id})(a) = h(a) 1 = (\text{id} \otimes h)(a)$ for all $a \in A$. This state is necessarily unique and it is faithful on $\mathcal{A}$. Then h may be called the Haar functional.

Woronowicz [9, §2] defines a *representation* of the compact matrix quantum group (A, u) on a finite dimensional vector space V as a linear mapping $t: V \rightarrow V \otimes A$ such that $(t \otimes \text{id}) \circ t = (\text{id} \otimes \Delta) \circ t$. If $t(v) = 0$ implies $v = 0$ then the representation is called *non-degenerate* and if $t(V) \subset V \otimes \mathcal{A}$ then the representation is called *smooth*. A smooth representation is non-degenerate iff $(\text{id} \otimes \varepsilon) \circ t = \text{id}$. Thus corepresentations of $\mathcal{A}$ on finite dimensional vector spaces, as defined in §1.3 of the present paper, correspond to nondegenerate smooth representations of (A, u) in [9].

As a consequence of the existence of the Haar functional, it is now shown (see [9, Theorem 5.2 and Prop. 3.2]) that nondegenerate smooth representations of (A, u) are unitarizable. Then we can conclude that the dense $*$ -algebra $\mathcal{A}$ of A is a CMQG algebra.

Conversely, if we start with a CMQG algebra $\mathcal{A}$ with fundamental corepresentation u as in Proposition 2.2 then it is possible to show the existence of a Haar functional h (cf. Theorem 2.8) without using C^* -algebras. Next a C^* -completion A can be made which uses the existence of the Haar functional (cf. §2.3). Then it is clear that the pair (A, u) is a compact matrix quantum group in the sense of Woronowicz. However, the C^* -algebra A as constructed from $\mathcal{A}$ in §2.3, is canonically determined by $\mathcal{A}$, but this is not the case with the compact matrix quantum

groups (A, u) of Woronowicz. In the present paper we have taken the completion of $\mathcal{A}$ with respect to the largest C^* -seminorm on $\mathcal{A}$, while, in [9], the norm induced by A on $\mathcal{A}$ may not be the largest C^* -seminorm on $\mathcal{A}$. In that case, the counit $\varepsilon: \mathcal{A} \rightarrow \mathbb{C}$ does not necessarily have a continuous extension to a linear functional on A (cf. [9, second Remark to Prop. 1.8]).

In his paper [10], Woronowicz starts with a compact matrix quantum group (A, u) in the sense of [9], then constructs out of its finite dimensional unitary representations a so-called complete concrete monoidal W^* -category (see [10, Theorem 1.2]) and next constructs from any such category a compact matrix quantum group (B, u) . Then B is not necessarily isomorphic to A , but B is obtained canonically from the CMQG algebra $\mathcal{A}$ generated by the entries of u , as in §2.3.

Both in [9] and in the present paper there is a similar key result [9, last statement of Theorem 5.4] resp. Proposition 2.7. We got the idea of the statement and proof of Proposition 2.7 from [9], but in the present paper, different from [9], the positivity and faithfulness of the Haar functional on $\mathcal{A}$ is a corollary rather than a prerequisite. As soon as these matters are settled, the quantum Schur orthogonality relations and the modular property of the Haar functional can be obtained in the same way in both papers, cf. [9, pp. 653–656] resp. §2.4 in the present paper.

(b) *Woronowicz [12] and S. Wang [8]*

Woronowicz [12] defines a *compact quantum group* as a pair (A, Δ) , where A is a separable unital C^* -algebra and $\Delta: A \rightarrow A \otimes A$ is a C^* -homomorphism, such that the following properties hold.

- 1) $(\Delta \otimes \text{id}) \circ \Delta = (\text{id} \otimes \Delta) \circ \Delta$.
- 2) $\text{Span}\{(b \otimes 1) \Delta(c) \mid b, c \in A\}$ and $\text{Span}\{(1 \otimes b) \Delta(c) \mid b, c \in A\}$ are dense subspaces of $A \otimes A$.

In particular, if (A, u) is a compact matrix pseudogroup as defined in [9] and if Δ is the corresponding comultiplication then (A, Δ) is a compact quantum group as just defined. Conversely, it is shown in [12] that, if (A, Δ) is a compact quantum group and if $\mathcal{A}$ is the set of all linear combinations of matrix elements of finite dimensional unitary representations of (A, Δ) then $\mathcal{A}$ is a dense $*$ -subalgebra of A and $\mathcal{A}$ is a Hopf $*$ -algebra. The existence of a Haar functional is also shown. It is observed that the representation theory as developed in [9] can be formulated in a similar way for compact quantum groups.

It is pointed out in Wang [8, Remark 2.2] that the results of [12] remain true if separability of the C^* -algebra A is no longer required, but if it is assumed instead that the C^* -algebra A has a faithful state. This observation would imply that a compact quantum group (A, Δ) in the sense of Wang gives rise to a CQG algebra $\mathcal{A}$ ($\mathcal{A}$ being constructed from A as in the previous paragraph), and that conversely each CQG algebra $\mathcal{A}$ would give rise to a compact quantum group (A, Δ) (A being completion of $\mathcal{A}$ with respect to maximal C^* -seminorm).

(c) *Effros & Ruan* [4]

In different terminology, CQG algebras were earlier introduced by Effros & Ruan [4]. They defined these algebras as cosemisimple Hopf algebras with a so-called standard $*$ -operation and they called these structures *discrete quantum groups*. This name was motivated by the fact that special examples of these algebras are provided by the group algebra of a discrete group, while the name CQG algebra comes from the class of examples, where we deal with the algebra of representative functions on a compact group. In the final section of [4] the authors define a *compact quantum group* as a natural generalization of the compact matrix quantum groups defined in [9]. Their definition involves a unital C^* -algebra A with a dense unital $*$ -subalgebra $\mathcal{A}$, where $\mathcal{A}$ is a CQG-algebra (in the terminology of the present paper) and the comultiplication on $\mathcal{A}$ extends continuously to A . Conversely, they show that a CQG algebra $\mathcal{A}$ gives rise to a compact matrix quantum group according to their definition. This involves a C^* -completion, for which a Haar functional h on $\mathcal{A}$ is needed. This Haar functional is obtained in a way very different from the method in the present paper. The authors first show the existence of a left Haar functional ϕ on a certain subspace of the linear dual of $\mathcal{A}$. Then h is constructed in terms of ϕ .

References

- [1] E. Abe, *Hopf algebras*, Cambridge University Press, 1980.
- [2] W. Arveson, *An invitation to C^* -algebra*, Springer, 1976.
- [3] M. S. Dijkhuizen, *On compact quantum groups and quantum homogeneous spaces*, Dissertation, University of Amsterdam, 1994.
- [4] E. G. Effros & Z.-J. Ruan, *Discrete quantum groups, I. The Haar measure*, preprint, 1993.
- [5] M. Hazewinkel, *Formal groups and applications*, Academic Press, 1978.
- [6] M. E. Sweedler, *Hopf algebras*, Benjamin, 1969.
- [7] E. C. Titchmarsh, *The theory of functions*, Oxford University Press, second ed., 1939.
- [8] S. Z. Wang, *General constructions of compact quantum groups*, preprint, 1993.
- [9] S. L. Woronowicz, *Compact matrix pseudogroups*, Comm. Math. Phys. 111 (1987), 613–685.
- [10] S. L. Woronowicz, *Tannaka-Krein duality for compact matrix pseudogroups. Twisted $SU(N)$ groups*, Invent. Math. 93 (1988), 35–76.
- [11] S. L. Woronowicz, *A remark on compact matrix quantum groups*, Lett. Math. Phys. 21 (1991), 35–39.
- [12] S. L. Woronowicz, *Compact quantum groups*, preprint, version August 1, 1993.